

2004

Evidence of red-cockaded woodpecker nestling displacement by southern flying squirrels.

Brent Burt

Stephen F Austin State University, dbburt@sfasu.edu

James R. McCormick

Davy Crockett National Forest, U.S.D.A. Forest Service, Rt. 1, Box 55FS, Kennard, TX 75847, jamesmccormick@fs.fed.us

Richard N. Conner

Southern Research Station, 506 Hayter, Street, Nacogdoches, TX 75965

Daniel Saenz

Wildlife Habitat and Silviculture Laboratory, Southern Research Station, U.S.D.A. Forest Service, Nacogdoches, TX 75962

Follow this and additional works at: <http://scholarworks.sfasu.edu/biology>



Part of the [Biology Commons](#)

[Tell us](#) how this article helped you.

Recommended Citation

McCormick, J. R., R. N. Conner, D. Saenz and D. B. Burt. 2004. Evidence of red-cockaded woodpecker nestling displacement by southern flying squirrels. *Bulletin of the Texas Ornithological Society* 37:8-9.

This Article is brought to you for free and open access by the Biology at SFA ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SFA ScholarWorks. For more information, please contact cdsscholarworks@sfasu.edu.

ILLiad Request Printout

Transaction Number: 17435
Username: burtdonal Name: Brent Burt
ISSN/ISBN: 0040-4543
NotWantedAfter: 03/10/2010
Accept Non English: No
Accept Alternate Edition: No
Request Type: Article - Article

Loan Information

LoanAuthor:
LoanTitle:
LoanPublisher:
LoanPlace:
LoanDate:
LoanEdition:
NotWantedAfter: 03/10/2010

Article Information

PhotoJournalTitle: Bulletin of the Texas Ornithological Society
PhotoJournalVolume: 37
PhotoJournalIssue:
Month:
Year: 2004
Pages: 8-9
Article Author: McCormick, J. R., R. N. Conner, D. Saenz and D. B. Burt.
Article Title: Evidence of red-cockaded woodpecker nestling displacement by southern flyin

Citation Information

Cited In:
Cited Title:
Cited Date:
Cited Volume:
Cited Pages:

dbburt@sfasu.edu

OCLC Information

ILL Number:
OCLC Number:
Lending String:
Original Loan Author:
Original Loan Title:
Old Journal Title:
Call Number: QL 671 .T42
Location:

Notes

2/8/2006 10:40:48 AM burtdonal QL671 .T42x Need in pdf format

- MONTGOMERY, T. H. 1907. The English Sparrow in Texas. *Auk* 24:341.
- OBERHOLSER, H. C. (n.d.). Typescript of *The Bird Life of Texas* (microfilm): *Passer domesticus*, pp. 8161–8192. Center for American History, University of Texas at Austin.
- PHILLIPS, J. C. 1915. Notes on American and old world English sparrows. *Auk* 32:51–59.
- "PROVO". 1895. Small birds and insects. *Forest and Stream* 45(12):248.
- RAGSDALE, G. H. 1892. Turn the rascals out. *Gainesville Hesperian*, 17 November 1892, p. 2.
- ROBBINS, C. S. 1973. Introduction, spread, and abundance of the House Sparrow in North America, pp. 3–9. In: S. Charles Kendeigh, A symposium on the House Sparrow (*Passer domesticus*) and European Tree Sparrow (*P. montanus*) in North America. AOU Monograph No. 14.
- SINGLEY, J. A. 1893. Texas birds: Notes on the birds of Galveston Island. Fourth Annual Report, Geological Survey of Texas for 1892, pp. 355–363.
- STRECKER, J. K. 1927. Notes on the ornithology of McLennan County, Texas. Baylor University Museum, Special Bulletin No. 1.

EVIDENCE OF RED-COCKADED WOODPECKER NESTLING DISPLACEMENT BY SOUTHERN FLYING SQUIRRELS

JAMES R. MCCORMICK¹, RICHARD N. CONNER^{2**}, DANIEL SAENZ², AND D. BRENT BURT³

¹Davy Crockett National Forest, U.S.D.A. Forest Service, Rt. 1, Box 55FS, Kennard, TX 75847, ²Wildlife Habitat and Silviculture Laboratory, Southern Research Station, U.S.D.A. Forest Service, Nacogdoches, TX 75962, and ³Department of Biology, Stephen F. Austin State University, Nacogdoches, TX 75962

ABSTRACT.—Red-cockaded Woodpeckers (*Picoides borealis*) are unique among woodpeckers in that they excavate their roost and nest cavities entirely within living pines (Ligon 1970). A number of secondary cavity nesters and other vertebrates are dependent on Red-cockaded Woodpeckers for the cavities they create (Rudolph et al. 1990, Loeb 1993, LaBranche and Walters 1994, Conner et al. 1997). Harlow and Lennartz (1983), Rudolph et al. (1990), Loeb (1993), and Conner et al. 1996, 1997) showed that southern flying squirrels (*Glaucomys volans*) in Texas and South Carolina were the most common occupants observed in the cavities other than Red-cockaded Woodpeckers and that these cavities were selected primarily on the basis of entrance size. While these studies showed that there is a propensity for squirrels to use unenlarged cavities, none have shown evidence of Red-cockaded Woodpeckers being killed by southern flying squirrels. Stabb et al. (1989) proposed that smaller birds may suffer mortality or disturbance from flying squirrels usurping the occupied cavities. Jackson (1978a), along with Hooper and Lennartz (1983), noted that southern flying squirrels have usurped Red-cockaded Woodpecker cavities without killing the birds. We report an instance of two 12- to 15-day-old nestlings found dead at the base of the nest tree and 3 flying squirrels inside the nest cavity.

While examining nest trees of resident groups of woodpeckers for presence of eggs and/or nestlings on the Angelina National Forest (62,423 ha; 31°15'N, 94°15'W) we inserted a small camera attached to a telescoping 15-m pole into each cavity chamber. On 26 May 1999 we observed two dead 12- to 15-day-old nestlings at the base of a nest tree. These nestlings were last observed alive in the cavity on 23 May 1999. Upon subsequent cavity inspection, we found the tree to be occupied by 3 southern flying squirrels. There was a well-developed, copious resin flow surrounding the cavity, but it did not deter the squirrels from entering the cavity, which is consistent with findings in other studies (Rudolph et al. 1990, Loeb 1993).

While it is conceivable that a predator such as a rat snake (*Elaphe obsoleta*) could have ejected the nestlings and the flying squirrels moved in after the cavity was unoccupied, certain evidence suggests that this was not the case. The resin barrier on this particular tree was well established with clear fresh resin 0.5 m above the cavity and 1.5 m below the cavity. Thus, it is unlikely that a snake or any other small terrestrial vertebrate could have gained access to the cavity contents. Dennis (1971) and Jackson (1978b) documented two instances of *E. obsoleta* gaining access

¹E-mail: jamesmccormick@fs.fed.us

²Maintained in cooperation with the Arthur Temple College of Forestry, Stephen F. Austin State University, Nacogdoches, TX.

^{**}Corresponding author: Richard N. Conner, Southern Research Station, 506 Hayter Street, Nacogdoches, TX 75965 Phone: (936) 569-7981.

to cavities occupied by flying squirrels or Red-cockaded Woodpecker nestlings, but the resin barriers on those cavities were compromised. Other study results (Jackson 1974, Rudolph et al. 1990) have supported the hypothesis that pine boles coated with fresh, sticky resin reduce the access that rat snakes have to cavities. Had a rat snake entered the cavity, it is almost certain that it would have eaten the nestlings. Given the fact that these birds were found 20 cm from the base of the nest tree, we suggest that the flying squirrels ejected the nestlings and usurped the cavity from the woodpeckers. Because there were no bite marks on either nestling and all adult woodpeckers were still present, we suggest that the cavity was usurped without attempting predation on the nestlings or adults.

LITERATURE CITED

- CONNER, R. N., D. C. RUDOLPH, D. SAENZ, AND R. R. SCHAEFER. 1996. Red-cockaded Woodpecker nesting success, forest structure, and southern flying squirrels in Texas. *Wilson Bulletin* 108: 697-711.
- . 1997. Species using Red-cockaded Woodpecker cavities in eastern Texas. *Bulletin of the Texas Ornithological Society* 30: 11-16.
- DENNIS, J. V. 1971. Utilization of pine resin by the Red-cockaded Woodpecker and its' effectiveness in protecting roosting and nesting sites, pp. 78-86 *In* R. L. Thompson, ed., *The ecology and management of the Red-cockaded Woodpecker*. Tallahassee: Bureau of Sport Fisheries and Wildlife, U. S. Department of the Interior and Tall Timbers Research Station.
- HARLOW, R. F. AND M. R. LENNARTZ. 1983. Interspecific competition for Red-cockaded Woodpecker cavities during the nesting season in South Carolina, pp. 41-43 D. A. Wood, ed., *In* Red-cockaded Woodpecker symposium II. Tallahassee: Florida Game and Freshwater Commission.
- HOOPER, R. G. AND M. R. LENNARTZ. 1983. Roosting behavior of Red-cockaded Woodpecker clans with insufficient cavities. *Journal of Field Ornithology* 54: 72-76.
- JACKSON, J. A. 1974. Gray rat snakes versus Red-cockaded Woodpecker: predator-prey adaptations. *Auk* 91: 342-347.
- . 1978a. Competition for cavities and Red-cockaded Woodpecker management, pp. 103-112 *In* S. A. Temple, ed., *Endangered Birds, Management Techniques for Preserving Threatened Species*. Madison: University of Wisconsin Press.
- . 1978b. Predation by a gray rat snake on Red-cockaded Woodpecker nestlings. *Bird-Banding* 49: 187-188.
- LABRANCHE, M. S. AND J. R. WALTERS. 1994. Patterns of mortality in nests of Red-cockaded Woodpeckers in the sandhills of southcentral North Carolina. *Wilson Bulletin* 106: 258-271.
- LIGON, J. D. 1970. Behavior and breeding biology of the Red-cockaded Woodpecker. *Auk* 87: 255-278.
- LOEB, S. C. 1993. Use and selection of Red-cockaded Woodpecker cavities by southern flying squirrels. *Journal of Wildlife Management* 57: 329-335.
- RUDOLPH, D. C., R. N. CONNER, AND J. TURNER. 1990. Competition for Red-cockaded Woodpecker roost and nest cavities: effects of resin age and entrance diameter. *Wilson Bulletin* 102: 23-36.
- STABB, M. A., M. E. GARTSHORE, AND P. L. AIRD. 1989. Interactions of southern flying squirrels *Glaucomys volans*, and cavity nesting birds. *Canadian Field-Naturalist* 103: 401-403.

DISTRIBUTION OF THE SUBSPECIES OF GREAT HORNED OWLS IN TEXAS

ROBERT W. DICKERMAN¹

*Museum of Southwestern Biology, University of New Mexico
Albuquerque, New Mexico 87131*

ABSTRACT.—The ranges of two subspecies of Great Horned Owls (*Bubo virginianus virginianus* and *B. v. pallescens*) in Texas were determined by comparison of 157 specimens from Texas and smaller series of each subspecies from east and west of Texas respectively, with the same three specimens (pale, medium and dark). Six color characters were used. The values of these were summed, and mapped. The range of nominate *virginianus* was found to be closely dependent on the distribution of hardwood, pine and oak dominated wooded areas in Texas.

The Great Horned Owl, *Bubo virginianus*, is an abundant, widespread, and successful avian predator in diverse habitats. Two distinctive subspecies occur in Texas. In the wooded areas of the east is the dark, heavily barred, richly ochaceous nominate subspecies, *B. v. virginianus*, while the paler, more xeric adapted subspecies, *B. v. pallescens*, (type locality: Watson Ranch, on the Medina River, 18 miles southwest of San

¹E-mail: bobdickm@unm.edu