

2009

# A new whole-stand model for unmanaged loblolly and slash pine plantations in east Texas

Dean W. Coble

*Arthur Temple College of Forestry and Agriculture, Stephen F. Austin State University, dcoble@sfasu.edu*

Follow this and additional works at: <http://scholarworks.sfasu.edu/forestry>



Part of the [Forest Sciences Commons](#)

Tell us how this article helped you.

---

## Recommended Citation

Coble, Dean W., "A new whole-stand model for unmanaged loblolly and slash pine plantations in east Texas" (2009). *Faculty Publications*. Paper 177.

<http://scholarworks.sfasu.edu/forestry/177>

This Article is brought to you for free and open access by the Forestry at SFA ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SFA ScholarWorks. For more information, please contact [cdsscholarworks@sfasu.edu](mailto:cdsscholarworks@sfasu.edu).

# A New Whole-Stand Model for Unmanaged Loblolly and Slash Pine Plantations in East Texas

Dean W. Coble

ABSTRACT

A new compatible whole-stand growth-and-yield model to predict total tree cubic-foot volume per acre yield (outside and inside bark) was developed for unmanaged loblolly pine (*Pinus taeda*) and slash pine (*Pinus elliottii*) plantations in East Texas. This model was compared with the noncompatible whole-stand model of Lenhart (Lenhart, 1996, Total and partial stand-level yield prediction for loblolly and slash pine plantations in east Texas, *South. J. Appl. For.* 20(1):36–41) and the Lenhart (1996) model refit to current data. For the two species, all three models were evaluated with independent observed data. The model developed in this study outperformed both Lenhart models in prediction of future yield and basal area per acre for all age classes combined and by 5-year age classes. The Lenhart models consistently overestimated yield and basal area per acre. All three models predicted surviving trees per acre similarly. An example is also provided to show users how to use the new whole-stand model.

**Keywords:** *Pinus taeda*, *Pinus elliottii*, compatible growth-and-yield models, Schumacher yield model

Whole-stand growth-and-yield models predict future yields as a function of stand-level attributes, such as site index (SI), age, and stand density (Avery and Burkhart 2002). MacKinney et al. (1937), MacKinney and Chaiten (1939), and Schumacher (1939) developed the earliest whole-stand growth-and-yield models. Clutter (1963) and Sullivan and Clutter (1972) used a Schumacher yield function as the basis for their compatible growth-and-yield model for loblolly pine stands. Buckman (1962) also developed a compatible growth-and-yield model for red pine stands in the Lake States. In a compatible growth-and-yield model, the yield function can be found by mathematically integrating the growth function. Subsequently, others used this concept to develop whole-stand growth-and-yield models for loblolly (Burkhart et al. 1972, 1985, Amateis et al. 1986, Ledbetter et al. 1986, Borders et al. 1990, Harrison and Borders 1996, Ochi and Cao 2003, Borders et al. 2004) and slash (Pienaar and Harrison 1989, Martin et al. 1999) pine plantations.

Lenhart (1996) developed a noncompatible whole-stand growth-and-yield model exclusively with data from unmanaged loblolly and slash pine plantations in East Texas. He used a Schumacher-type function to predict future yields for different merchantability standards (Amateis et al. 1986). However, his data represented relatively young plantations (average age, 10 years; range, 5–24 years) commonly established in the early 1980s. These plantations are still found in East Texas, but they are currently older than those represented by Lenhart (1996).

The objective of this study was to develop new whole-stand growth-and-yield models for unmanaged loblolly and slash pine plantations in East Texas. The intent was to improve on Lenhart (1996) by using new equations and his original data set with an

additional 15 years of new measurements from older plantations to forecast future yields.

## Data Description

This study used 987 observations from 173 remeasured permanent plots located in East Texas loblolly pine plantations (Table 1). From the total 987 observations, approximately 10% ( $n = 103$  observations from 18 permanent plots) were randomly selected and removed from the data set used for model fitting and reserved for model evaluation. Thus, a total of 884 loblolly pine observations from 155 permanent plots were used for model fitting. For slash pine, a total of 430 observations from 78 permanent plots were used (Table 2). Ten percent ( $n = 40$  observations from 8 permanent plots) were removed from the model fitting data set and reserved for model evaluation, leaving 390 observations from 70 plots for model fitting.

The 173 loblolly pine and 78 slash pine permanent plots are part of the East Texas Pine Plantation Research Project (ETPPRP; Lenhart et al. 1985), which covers 22 counties across East Texas. Generally, the counties are located within the rectangle from 30 to 35° north latitude and 93 to 96° west longitude. Each plot consists of two adjacent subplots approximately 0.25 ac in size (100 × 100-ft) separated by a 60-ft buffer. Within a subplot, dbh (measured at 4.5 ft above the groundline), total height, and the survival status (live or dead) were monitored for each planted loblolly pine tree over a 27-year period (note: Lenhart (1996) used the same data set, but the plots had only been monitored over a 12-year period). Plots were remeasured on fixed, 3-year intervals. Data from only one subplot (the development subplot) were used in this study. For each tree, dbh and total height were used to estimate total tree (from the stump to the tree tip) cubic-foot volume (outside bark (ob) and inside bark

Received January 15, 2008; accepted April 9, 2008.

Dean Coble (dcoble@sfasu.edu), Arthur Temple College of Forestry and Agriculture, Stephen F. Austin State University, Box 6109 SFA Station, Nacogdoches, TX 75962. The author thanks Temple-Inland, International Paper, and Stephen F. Austin State University for funding this study through the East Texas Pine Plantation Research Project. The author is indebted to Dr. J. David Lenhart for creating the ETPPRP as well as all the people that helped collect the data over the years. The author also thanks Dehai Zhao, Chris Cieszewski, Quang Cao, Mike Strub, and Bruce Borders for their helpful comments at the 2007 Southern Mensurationists Meeting as well as three anonymous reviewers.

Copyright © 2009 by the Society of American Foresters.

**Table 1. Observed stand characteristics for East Texas unmanaged loblolly pine plantation data sets for whole-stand model.**

Variables	Model development data set (n = 884 observations from 155 plots)				Model evaluation data set (n = 103 observations from 18 plots)			
	Mean	SD	Minimum	Maximum	Mean	SD	Minimum	Maximum
A	14.0	6.6	2.0	37.0	14.2	6.9	3.0	32.0
H	44.7	19.3	5.0	93.0	45.1	20.8	7.0	93.0
SI	72.1	10.0	33.0	96.0	72.5	9.9	53.0	88.0
TPA	455.5	147.7	83.0	1,002.0	437.8	147.5	170.0	858.0
BA	89.7	52.2	0.5	220.7	85.9	54.6	0.7	215.3
Dq	5.8	2.4	0.1	12.9	5.7	2.5	0.3	11.5
V(ob)	1,870.6	1,523.7	0.1	6,607.7	1,833.3	1,596.7	0.4	6,135.2
V(ib)	1,545.8	1,257.4	0.1	5,450.0	1,514.8	1,317.6	0.3	5,063.7

A, plantation age (total yr); H, average height of dominant and codominant trees (ft); SI, site index (base age, 25 yr); TPA, trees per acre; BA, loblolly pine basal area per acre (BAPA, ft<sup>2</sup>); Dq = quadratic mean diameter (inches); V = total tree cubic-foot volume per acre, SD, standard deviation.

**Table 2. Observed stand characteristics for East Texas unmanaged slash pine plantation data sets for whole-stand model.**

Variables	Model development data set (n = 390 observations from 70 plots)				Model evaluation data set (n = 40 observations from 8 plots)			
	Mean	SD	Minimum	Maximum	Mean	SD	Minimum	Maximum
A	13.4	6.2	2.0	31.0	13.2	6.1	3.0	27.0
H	42.7	19.3	4.0	91.0	41.6	19.8	8.0	87.0
SI	74.7	10.0	36.0	101.0	72.8	5.0	65.0	81.0
TPA	374.9	173.4	61.0	1,002.0	369.0	182.8	70.0	671.0
BA	64.7	40.7	0.5	162.3	54.4	37.6	0.8	120.5
Dq	5.7	2.5	0.1	11.5	5.5	2.8	0.3	11.7
V(ob)	1,500.6	1,267.7	0.1	5,430.1	1,260.6	1,198.9	1.3	4,453.8
V(ib)	1,139.0	1,000.0	0.1	4,408.3	958.2	957.4	0.7	3,627.1

A, plantation age (total yr); H, average height of dominant and codominant trees (ft); SI, site index (base age, 25 yr); TPA, trees per acre; BA, loblolly pine basal area per acre (BAPA, ft<sup>2</sup>); Dq, quadratic mean diameter (inches); V, total tree cubic-foot volume per acre; SD, standard deviation.

(ib)). The cubic-foot volume equations of Coble and Hilpp (2006) were used for loblolly pine, and the cubic-foot volume equations of Lenhart et al. (1987) were used for slash pine. Dominant height was determined by averaging the heights of the tallest 10 trees on a subplot (which approximately represents the tallest 40 trees per acre) that were free of damage, forks, and stem fusiform rust (*Cronartium quercuum* [Berk.] Miyabe ex Shirai f. sp. *Fusiforme*).

### The New Whole-stand Model

The Schumacher (1939) model was used to predict future total tree cubic-foot volume per acre ( $V_2$ ) as a function of future dominant height, plantation age, and basal area per acre:

$$\ln V_2 = a_0 + a_1 \ln H_2 + a_2 \frac{1}{A_2} + a_3 \ln B_2, \quad (1)$$

where  $H_2$  = future average height (feet) of dominant and codominant trees;  $A_2$  = future plantation age (years);  $B_2$  = future basal area (squared feet) per acre;  $a_i$  = parameter estimates; ln = natural logarithm function.

Future basal area per acre ( $B_2$ ) and future dominant height ( $H_2$ ) must be known before Equation 1 can be used. The model of Borders et al. (2004) was used to predict current basal area per acre (squared feet,  $B_1$ ):

$$\ln B_1 = b_0 + b_1 \frac{1}{A_1} + b_2 \ln N_1 + b_3 \ln H_1 + b_4 \ln N_1 \frac{1}{A_1} + b_5 \ln H_1 \frac{1}{A_1}, \quad (2)$$

where  $A_1$  = current plantation age (years);  $N_1$  = current trees per acre;  $H_1$  = current average height (feet) of dominant and codominant trees;  $b_i$  = parameter estimates.

To predict future basal area per acre ( $B_2$ ), Borders et al. (2004) derived a basal area projection equation from Equation 2 by isolating the  $b_1$  term in Equation 2:

$$\ln B_2 = c_0 + \frac{A_1}{A_2} \left( \ln B_1 - c_0 - c_1 \ln N_1 - c_2 \ln H_1 - c_3 \frac{\ln N_1}{A_1} - c_4 \frac{\ln H_1}{A_1} \right) + c_1 \ln N_2 + c_2 \ln H_2 + c_3 \frac{\ln N_2}{A_2} + c_4 \frac{\ln H_2}{A_2}, \quad (3)$$

where  $N_2$  = future trees per acre;  $c_i$  = parameter estimates; and all other variables are as defined previously.

A Chapman-Richards model (Coble and Lee 2006) was used to predict future dominant height ( $H_2$ ):

$$H_2 = H_1 \left( \frac{1 - e^{-d_1 A_2}}{1 - e^{-d_1 A_1}} \right)^{d_2}, \quad (4)$$

where  $d_i$  = parameter estimates;  $e$  = exponential function; and all other variables are as defined previously.

Finally, a negative-exponential survival model (Zhao et al. 2007, Equation 32) was used to predict future surviving trees per acre ( $N_2$ ), which is an independent variable in Equation 3:

$$N_2 = N_1 e^{k \cdot \text{SI}(A_2 - A_1)}, \quad (5)$$

**Table 3. Parameter estimates and fit statistics of East Texas loblolly pine plantation predictive equations for ob future total tree cubic-foot volume per acre ( $\ln V_2$ ), current basal area per acre ( $\text{ft}^2$ ,  $\ln B_1$ ), future basal area per acre ( $\text{ft}^2$ ,  $\ln B_2$ ), future dominant height ( $\text{ft}$ ,  $H_2$ ), and future surviving trees per acre ( $N_2$ ).**

Equation	Parameter	Parameter estimate	Standard error	Pr(parameter = 0)	$R^2$	RMSE
1 ( $\ln V_2$ )	$a_0$	-1.53354445	0.0394	<0.0001	0.998	0.0456
	$a_1$	1.16418329	0.0116	<0.0001		
	$a_2$	0.26335955	0.0946	0.0055		
	$a_3$	0.97534638	0.0049	<0.0001		
	$b_0$	-5.12109968	0.2527	<0.0001		
2 ( $\ln B_1$ )	$b_1$	-14.89666626	1.9164	<0.0001	0.968	0.2169
	$b_2$	0.55230267	0.0317	<0.0001		
	$b_3$	1.55304841	0.0310	<0.0001		
	$b_4$	-0.14111514	0.2859	0.6217		
	$b_5$	5.27278019	0.1659	<0.0001		
3 ( $\ln B_2$ )	$c_0$	-3.19205665	0.3718	<0.0001	0.947	0.1425
	$c_1$	0.58186175	0.0485	<0.0001		
	$c_2$	1.09433140	0.0436	<0.0001		
	$c_3$	0.74064246	1.1107	0.5050		
	$c_4$	5.17072095	0.1886	<0.0001		
4 ( $H_2$ )	$d_1$	0.07811158	0.0039	<0.0001	0.943	4.1515
	$d_2$	1.48389930	0.0411	<0.0001		
5 ( $N_2$ )	$k$	-0.00021308	7.495E-6	<0.0001	0.973	23.5832

RMSE, root mean square error.

**Table 4. Parameter estimates and fit statistics of East Texas loblolly pine plantation predictive equations for ib future total tree cubic-foot volume per acre ( $\ln V_2$ ), current basal area per acre ( $\text{ft}^2$ ,  $\ln B_1$ ), future basal area per acre ( $\text{ft}^2$ ,  $\ln B_2$ ), future dominant height ( $\text{ft}$ ,  $H_2$ ), and future surviving trees per acre ( $N_2$ ).**

Equation	Parameter	Parameter estimate	Standard error	Pr(parameter = 0)	$R^2$	RMSE
1 ( $\ln V_2$ )	$a_0$	-1.70171933	0.0394	<0.0001	0.998	0.0454
	$a_1$	1.15943023	0.0116	<0.0001		
	$a_2$	0.27127090	0.0942	0.0041		
	$a_3$	0.97456387	0.0049	<0.0001		
	$b_0$	-5.11826231	0.2527	<0.0001		
2 ( $\ln B_1$ )	$b_1$	-14.91102867	1.9167	<0.0001	0.968	0.2169
	$b_2$	0.55188370	0.0317	<0.0001		
	$b_3$	1.55275343	0.0310	<0.0001		
	$b_4$	-0.14150303	0.2860	0.6208		
	$b_5$	5.28023138	0.1659	<0.0001		
3 ( $\ln B_2$ )	$c_0$	-3.19383800	0.3718	<0.0001	0.946	0.1425
	$c_1$	0.58176619	0.0485	<0.0001		
	$c_2$	1.09492902	0.0436	<0.0001		
	$c_3$	0.73979043	1.1105	0.5055		
	$c_4$	5.16835658	0.1886	<0.0001		
4 ( $H_2$ )	$d_1$	0.07811170	0.0039	<0.0001	0.943	4.1515
	$d_2$	1.48386038	0.0411	<0.0001		
5 ( $N_2$ )	$k$	-0.00021310	7.495E-6	<0.0001	0.973	23.5832

where SI = site index in feet (index age, 25 years);  $k$  = parameter estimate; and all other variables are as defined previously.

Equations 1–5 were simultaneously fit to the fitting data set using the MODEL Procedure in SAS/ETS (SAS Institute, Inc., 2004) to obtain parameter estimates for model evaluation. After model evaluation was complete, the fitting and evaluation data sets were combined into one data set. Equations 1–5 were then simultaneously fit to the combined data to obtain final parameter estimates (Tables 3–6). In all cases, seemingly unrelated regression was used to account for correlation across the equations (Borders 1989, Robinson 2004).

## Model Evaluation

The new whole-stand model's equations for future yield (Equation 1), basal area per acre (Equation 3), and surviving trees per acre (Equation 5) were evaluated with the 10% evaluation data set. Lenhart's (1996) equations for  $\ln V_2$ ,  $\ln B_2$ , and  $N_2$  were

also evaluated with the 10% evaluation data set to determine if the new model was an improvement over Lenhart (1996). Lenhart's equations for  $\ln V_2$ ,  $\ln B_2$ , and  $N_2$  were also fit to the current development data set to provide a current comparison with the new model (hereafter called modified Lenhart). This third comparison allows for a separate evaluation of model form and data age in Lenhart (1996). In Lenhart's model,  $\ln B_2$  is not predicted directly, but rather natural logarithm of future quadratic mean diameter ( $\ln Dq_2$ ). However,  $\ln B_2$  was derived from his  $N_2$  and  $\ln Dq_2$  equations for purposes of comparison.

All comparisons were made in original units rather than natural logarithm units thereby making the comparisons more meaningful to the reader. Thus,  $V_2$ ,  $B_2$ , and  $N_2$  for all three models (new model; Lenhart 1996, modified Lenhart) were evaluated with the 10% evaluation data set.

Evaluation of all three models was performed at two levels of resolution: (1) all age classes and (2) 5-year age classes (from 5 to 30

**Table 5. Parameter estimates and fit statistics of East Texas slash pine plantation predictive equations for ob future total tree cubic-foot volume per acre (ln  $V_2$ ), current basal area per acre (ft<sup>2</sup>, ln  $B_1$ ), future basal area per acre (ft<sup>2</sup>, ln  $B_2$ ), future dominant height (ft,  $H_2$ ), and future surviving trees per acre ( $N_2$ ).**

Equation	Parameter	Parameter estimate	Standard error	Pr(parameter = 0)	$R^2$	RMSE
1 (ln $V_2$ )	$a_0$	-1.23873624	0.0483	<0.0001	0.999	0.0387
	$a_1$	1.13402680	0.0132	<0.0001		
	$a_2$	0.39260901	0.1199	0.0011		
	$a_3$	0.96958127	0.0048	<0.0001		
	$b_0$	-6.23132969	0.3001	<0.0001		
2 (ln $B_1$ )	$b_1$	-10.93367473	2.5930	<0.0001	0.966	0.2115
	$b_2$	0.64518403	0.0361	<0.0001		
	$b_3$	1.64301972	0.0363	<0.0001		
	$b_4$	-0.08801420	0.3733	0.8137		
	$b_5$	4.55471494	0.2623	<0.0001		
	$c_0$	-2.80921145	0.4280	<0.0001		
3 (ln $B_2$ )	$c_1$	0.63062849	0.0463	<0.0001	0.971	0.1106
	$c_2$	0.97003672	0.0553	<0.0001		
	$c_3$	2.46131790	1.1700	0.0360		
	$c_4$	2.44542174	0.2660	<0.0001		
	$d_1$	0.05225618	0.0051	<0.0001		
4 ( $H_2$ )	$d_2$	1.25939432	0.0468	<0.0001	0.955	3.7808
	$k$	-0.00030932	0.000013	<0.0001		
5 ( $N_2$ )					0.979	24.7091

**Table 6. Parameter estimates and fit statistics of East Texas slash pine plantation predictive equations for ib future total tree cubic-foot volume per acre (ln  $V_2$ ), current basal area per acre (ft<sup>2</sup>, ln  $B_1$ ), future basal area per acre (ft<sup>2</sup>, ln  $B_2$ ), future dominant height (ft,  $H_2$ ), and future surviving trees per acre ( $N_2$ ).**

Equation	Parameter	Parameter estimate	Standard error	Pr (parameter = 0)	$R^2$	RMSE
1 (ln $V_2$ )	$a_0$	-2.24692686	0.0546	<0.0001	0.998	0.0450
	$a_1$	1.33647003	0.0149	<0.0001		
	$a_2$	0.41097117	0.1359	0.0026		
	$a_3$	0.95130442	0.0054	<0.0001		
	$b_0$	-6.22771268	0.2936	<0.0001		
2 (ln $B_1$ )	$b_1$	-11.29231165	2.5142	<0.0001	0.965	0.2115
	$b_2$	0.64797833	0.0352	<0.0001		
	$b_3$	1.64255519	0.0359	<0.0001		
	$b_4$	-0.00158467	0.3618	0.9965		
	$b_5$	4.45116276	0.2549	<0.0001		
	$c_0$	-2.90167368	0.4240	<0.0001		
3 (ln $B_2$ )	$c_1$	0.64055826	0.0460	<0.0001	0.971	0.1107
	$c_2$	0.97614432	0.0546	<0.0001		
	$c_3$	2.21070791	1.1552	0.0563		
	$c_4$	2.46616364	0.2641	<0.0001		
	$d_1$	0.05232592	0.0051	<0.0001		
4 ( $H_2$ )	$d_2$	1.26007790	0.0468	<0.0001	0.955	3.7808
	$k$	-0.00030950	0.000013	<0.0001		
5 ( $N_2$ )					0.979	24.7091

years). For the all age class evaluation, four criteria (Kozak and Smith 1993) were used:

- Mean Bias = Bias =  $\frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)}{n}$ ,
- Mean Percent bias = %Bias =  $\frac{\sum_{i=1}^n \left[ 100 \left( \frac{Y_i - \hat{Y}_i}{Y_i} \right) \right]}{n}$ ,
- Standard error of the estimate = SEE =  $\sqrt{\frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}{n - k}}$ ,
- Percent SEE = %SEE =  $\left( \frac{SEE}{\bar{Y}} \right) 100$ ,

where  $Y_i$  = observed  $V_2$ ,  $B_2$ , and  $N_2$  for observation  $i$ ;  $\hat{Y}_i$  = predicted  $V_2$ ,  $B_2$ , and  $N_2$  for observation  $i$ ;  $\bar{Y}$  = mean  $V_2$ ,  $B_2$ , and  $N_2$ ;  $n$  = number of observations;  $k$  = number of estimated parameters in equation ( $k = 4$  for Equation 1,  $k = 5$  for Equation 3,  $k = 1$  for Equation 5, and  $k = 4$  for all equations of Lenhart 1996, and modified Lenhart). Note that a negative mean bias value corresponds to an overprediction while a positive mean bias value corresponds to an underprediction.

For the 5-year age class evaluation, mean predicted and observed  $V_2$ ,  $B_2$ , and  $N_2$  were calculated for each 5-year age class. For all three models, mean bias (defined previously) was then calculated for  $V_2$ ,  $B_2$ , and  $N_2$  by 5-year age classes. Mean bias was next plotted over age class to further examine model prediction trends across the range of ages, where negative values represent overpredictions and positive values represent underpredictions.

**Table 7. Loblolly pine mean bias, mean %bias, SEE, %SEE, and number of samples (n) for both ob and ib predicted future yield ( $V_2$ ), future basal area per acre ( $B_2$ ), and future surviving trees per acre ( $N_2$ ) from three growth and yield models: this study (ETPPRP), Lenhart (1996), and Lenhart (1996) refit with current data (modified Lenhart).**

Equation, Component, and Model	n	Mean bias	Mean %bias	SEE	%SEE
<b>1 <math>V_2</math></b>					
Wood and bark (ob)					
ETPPRP	103	-26.97	-0.92	87.26	3.7
Lenhart (1996)	103	-800.29	-47.30	1,373.89	58.21
Modified Lenhart	103	-180.90	-24.23	603.55	25.57
Wood only (ib)					
ETPPRP	103	-22.00	-0.92	71.36	3.66
Lenhart (1996)	103	-671.20	-40.96	1,231.94	63.18
Modified Lenhart	103	-148.98	-24.17	497.06	25.49
<b>3 <math>B_2</math></b>					
ob and ib					
ETPPRP	103	-0.87	-2.87	10.73	10.41
Lenhart (1996)	103	-25.93	-29.73	44.18	42.89
Modified Lenhart	103	-5.98	-16.33	20.59	19.99
<b>5 <math>N_2</math></b>					
ob and ib					
ETPPRP	103	-2.08	-0.98	24.74	5.94
Lenhart (1996)	103	-6.31	-2.04	25.76	6.19
Modified Lenhart	103	-0.80	-0.73	25.20	6.05

## Results and Discussion

The parameter estimates for loblolly and slash pine (ob and ib) using the combined data set (fitting plus evaluation data) were reported for Equations 1–5 (Tables 3–6). All parameter estimates were significantly different from zero at the 0.01 significance level, except for the  $\ln N_1/A_1$  term in Equation 2 as well as the  $\ln N_2/A_2$  term in Equation 3. Although these two interaction terms were not significant, they were not removed from the final equations so not to alter the mathematical compatibility between the basal area prediction (Equation 2) and projection (Equation 3) equations. All  $R^2$  values for each individual equation exceeded 94%, and the  $R^2$  value for the yield model (Equation 1) exceeded 99%. Not surprisingly, the intercept term of the yield model (Equation 1) was the only parameter value that differed dramatically between ob and ib measurements within a species.

Predicted values for  $V_2$  and  $N_2$  from the new model were less than 1% of observed values for loblolly pine, while predicted values for  $B_2$  were less than 3% (Table 7). For slash pine, predicted values for  $V_2$  were also less than 1% and  $B_2$  were less than 3% of observed values (Table 8). However, the new model overpredicted  $N_2$  by about 9% (mean %bias = 8.97, Table 8). Lenhart (1996) overpredicted  $V_2$  by about 41–47%,  $B_2$  by about 30%, and  $N_2$  by about 2% for loblolly pine (Table 7). For slash pine, Lenhart (1996) overpredicted  $V_2$  by about 12%,  $B_2$  by about 14%, and  $N_2$  by about 7% (Table 8). Modified Lenhart overpredicted  $V_2$  by about 24% and  $B_2$  by about 16% for loblolly pine (Table 7). However, modified Lenhart predicted  $N_2$  within 1% of observed values for loblolly pine (Table 7). For slash pine, modified Lenhart overpredicted  $V_2$  by about 18%,  $B_2$  by about 15%, and  $N_2$  by about 7% (Table 8). For Lenhart's model, predictions of  $V_2$ ,  $B_2$ , and  $N_2$  for loblolly pine improved with the addition of the older plantation data; all mean percent bias values were lower for modified Lenhart compared with Lenhart (1996; Table 7). This trend did not hold for slash pine (Table 8), although both versions of Lenhart better predicted  $V_2$  and  $B_2$  (but not  $N_2$ ) for slash pine than loblolly pine (Tables 7 and 8).

However, even using the current data set (thereby eliminating age differences the data), modified Lenhart was unable to predict  $V_2$  and  $B_2$  for loblolly and slash pine as well as the new model in terms of mean percent bias. However, all three models were similar in predicting  $N_2$ . Furthermore, all values of %SEE for the new model were lower than those of Lenhart of either model (Tables 7 and 8). Thus, based on this, all age class evaluation (mean %bias and %SEE), the new model better predicts future yield and basal area per acre than Lenhart (1996), or modified Lenhart, while prediction of survival is similar between the three models.

For 5-year age classes, loblolly and slash pine (ob) mean bias values for  $V_2$  and  $B_2$  for the new model fell closest to the zero line compared with Lenhart (1996) or modified Lenhart, (Figures 1 and 2). In fact, all values for  $V_2$  and  $B_2$  were near zero, which means that the new model was not over- or underpredicting to any appreciable degree. Modified Lenhart increasingly overpredicted  $V_2$  and  $B_2$  for loblolly pine as plantation age class increases. Lenhart (1996) also overpredicted  $V_2$  and  $B_2$  for slash pine as plantation age class increases, although not as dramatically as for loblolly pine. The only exception was for the 25-year age class; Lenhart (1996) underpredicted this age class, and then dropped near or below the zero line for the 30-year age class. Lenhart (2008) performed similarly to Lenhart (1996), although its trend line was closer to the zero line than Lenhart (1996). This is indicative of the improvement from using current data; predictions were improved using data from older plantations, although they never surpassed those of the new model. Mean bias values for  $N_2$  were similar between the three models for loblolly and slash pine (Figure 3). For both species and all three models, survival for age classes less than 25 years old was predicted best, with the worst predictions for the oldest two age classes. Although the models predicted  $N_2$  similarly, the superior performance by the new model for the  $V_2$  and  $B_2$  predictions represents an improvement over Lenhart (1996, 2008). Note that the mean bias trends by age class for ib  $V_2$ ,  $B_2$ , and  $N_2$  were similar to those for ob (figures not presented).

**Table 8. Slash pine mean bias, mean %bias, SEE, %SEE, and number of samples (n) for both ob and ib predicted future yield ( $V_2$ ), future basal area per acre ( $B_2$ ), and future surviving trees per acre ( $N_2$ ) from three growth-and-yield models: this study (ETPPRP), Lenhart (1996), and Lenhart refit with current data (modified Lenhart).**

Equation, Component, and Model	n	Mean bias	Mean %Bias	SEE	%SEE
<b>1 <math>V_2</math></b>					
Wood and bark (ob)					
ETPPRP	40	8.56	-0.38	105.67	6.70
Lenhart (1996)	40	53.27	-12.98	519.46	32.95
Modified Lenhart	40	-92.17	-18.51	479.92	30.45
Wood only (ib)					
ETPPRP	40	5.09	-0.81	93.95	7.77
Lenhart (1996)	40	72.37	-12.09	431.39	35.66
Modified Lenhart	40	-65.63	-19.67	386.87	31.98
<b>3 <math>B_2</math></b>					
ob and ib					
ETPPRP	40	-1.52	-2.57	6.34	9.96
Lenhart (1996)	40	-3.90	-13.62	17.40	27.35
Modified Lenhart	40	-4.54	-15.28	17.26	27.12
<b>5 <math>N_2</math></b>					
ob and ib					
ETPPRP	40	-14.56	-8.97	33.90	10.23
Lenhart (1996)	40	-15.17	-7.47	38.05	11.49
Modified Lenhart	40	-13.55	-7.06	37.11	11.20




Figure 1. Mean bias of future yield ( $V_2$ ) for loblolly and slash pine (ob) from three growth and yield models: this study (ETPPRP), Lenhart (1996), and Lenhart refit with current data (modified Lenhart).




Figure 2. Mean bias of future basal area per acre ( $B_2$ ) for loblolly and slash pine (ob) from three growth-and-yield models: this study (ETPPRP), Lenhart (1996), and Lenhart refit with current data (modified Lenhart).




Figure 3. Mean bias of future surviving trees per acre ( $N_2$ ) for loblolly and slash pine (ob) three growth-and-yield models: this study (ETPPRP), Lenhart (1996), and Lenhart refit with current data (modified Lenhart).

In conclusion, the new model better predicted  $V_2$  and  $B_2$  versus the two Lenhart models. All three models predicted  $N_2$  similarly. Lenhart's model benefited from refitting to current data; modified Lenhart outperformed Lenhart (1996) in terms bias (mean %bias) and precision (%SEE). The plantation age in the data set used by

Lenhart (1996) averaged 10–11 years old versus 13–14 years old in this study (also, maximum plantation age was 24 years old versus 30+ years, respectively). Clearly, additional data from older plantations improved performance in the Lenhart model. However, modified Lenhart still failed to outperform the new model in terms

of bias and precision. I believe the new model in this study benefits from being compatible as well as fit with simultaneous estimation techniques, and these two characteristics make the new model preferable to Lenhart's model.

## Application

To show how to use the new model, I will illustrate how to calculate the ob cubic-foot yield at rotation age =  $A_2 = 25$  years for a loblolly pine plantation with SI = 70 ft (index age, 25 years). Let  $A_1 = 1$  year and  $N_1 =$  initial planting density = 605 trees per acre.

1. Calculate  $H_1$  from Equation 4 by setting  $H_1 =$  SI = 70,  $A_1 =$  Index age = 25,  $H_2 = H_1$ , and  $A_2 = A_1 = 1$ :

$$H_1 = 70 \left( \frac{1 - e^{-0.07811158 * 1}}{1 - e^{-0.07811158 * 25}} \right)^{1.48389930} = 1.886 \text{ feet.}$$

Note that you can also obtain this value from field measurements.

2. Calculate  $H_2$  from Equation 4 using the value for  $H_1$  from the previous equation:

$$H_2 = 1.886 \left( \frac{1 - e^{-0.07811158 * 25}}{1 - e^{-0.07811158 * 1}} \right)^{1.48389930} = 70.0 \text{ feet.}$$

3. Calculate  $N_2$  from Equation 5:

$$N_2 = 605 e^{-0.00021308 * 70(25-1)} = 422.95 \text{ tpa.}$$

4. Calculate  $B_1$  using Equation 2:

$$\begin{aligned} \ln B_1 &= -5.12109968 - 14.89666626 * \frac{1}{1} \\ &+ 0.55230267 * \ln(605) \\ &+ 1.55304841 * \ln(1.886) - 0.14111514 * \frac{\ln(605)}{1} \\ &+ 5.27278019 * \frac{\ln(1.886)}{1} \\ &= -13.053313. \end{aligned}$$

$$\text{So, } B_1 = e^{-13.053313} = 0.000002 \text{ ft}^2/\text{ac.}$$

5. Calculate  $B_2$  using Equation 3:

$$\begin{aligned} \ln B_2 &= 3.19205665 \\ &+ \frac{1}{25} \left( \ln(0.000002) + 3.19205665 - 0.58186175 * \ln(605) \right. \\ &\quad \left. - 1.09433140 * \ln(1.886) - 0.74064246 * \frac{\ln(605)}{1} \right. \\ &\quad \left. - 5.17072095 * \frac{\ln(1.886)}{1} \right) \\ &+ 0.58186175 * \ln(422.95) + 1.09433140 * \ln(70.0) \\ &+ 0.74064246 * \frac{\ln(422.95)}{25} \\ &+ 5.17072095 * \frac{\ln(70.0)}{25} = 5.13879. \end{aligned}$$

$$\text{So, } B_2 = e^{5.13879} = 170.49225 \text{ or } 170.49 \text{ ft}^2/\text{ac.}$$

6. Calculate  $V_2$  using Equation 1:

$$\begin{aligned} \ln V_2 &= -1.53354445 + 1.16418329 * \ln(70.0) \\ &+ 0.2633595 * \frac{1}{25} \\ &+ 0.97534638 * \ln(170.49225) \\ &= 8.43502. \end{aligned}$$

$$\text{So, } V_2 = e^{8.43502} = 4605.5604 \text{ or } 4,606 \text{ ft}^3/\text{ac.}$$

## Literature Cited

- AMATEIS, R.L., H.E. BURKHART, AND T.E. BURK. 1986. A ratio approach to predicting merchantable yields of unthinned loblolly pine plantations. *For. Sci.* 32(2):287-296.
- AVERY, T.E., AND H.E. BURKHART. 2002. *Forest measurements*, 5th Ed. McGraw-Hill, Inc., New York. 456 p.
- BORDERS, B.E. 1989. Systems of equations in forest stand modeling. *For. Sci.* 35(2):548-556.
- BORDERS, B.E., W.M. HARRISON, D.E. ADAMS, R.L. BAILEY, AND L.V. PIENAAR. 1990. *Yield prediction and growth projection for site-prepared loblolly pine plantations in the Carolinas, Georgia, Florida, and Alabama*. University of Georgia School of Forest Resources PMRC Res. Pap. 1990-2, Univ. of Georgia, Athens, GA. 65 p.
- BORDERS, B.E., W.M. HARRISON, F.Y. ZHANG, B.D. SHIVER, M. CLUTTER, C. CIESZEWSKI, AND R.F. DANIELS. 2004. *Growth and yield models for second rotation loblolly pine plantations in the Piedmont/Upper Coastal Plain and Lower Coastal Plain of the southeastern U.S.* University of Georgia School of Forest Resources PMRC Res. Pap., Univ. of Georgia, Athens, GA. 64 p.
- BUCKMAN, R.E. 1962. *Growth and yield of red pine in Minnesota*. US For. Serv. Tech. Bull. 1272. 50 p.
- BURKHART, H.E., R.C. PARKER, M.R. STRUB, AND R.G. ODERWALD. 1972. *Yields of old-field loblolly pine plantations*. VPI&SU FWS-3-72. 51 p.
- BURKHART, H.E., D.C. CLOEREN, AND R.L. AMATEIS. 1985. Yield relationships in unthinned loblolly pine plantations on cutover, site-prepared lands. *South. J. Appl. For.* 9(2):84-91.
- CLUTTER, J.L. 1963. Compatible growth and yield models for loblolly pine. *For. Sci.* 9(3):354-371.
- COBLE, D.W., AND Y.J. LEE. 2006. Use of a generalized sigmoid growth function to predict site index for unmanaged loblolly and slash pine plantations in east Texas. P. 291-295 in *Proc. of the 13th Biennial Southern Silvicultural Research Conf., 2006*, Connor, K.F. (ed.). US For. Serv. Gen. Tech. Rep. SRS-92.
- COBLE, D.W., AND K. HILPP. 2006. Compatible cubic-foot stem volume and upper-stem diameter equations for semi-intensive plantation grown loblolly pine trees in east Texas. *South. J. Appl. For.* 30(3):132-141.
- HARRISON, W.M., AND B.E. BORDERS. 1996. *Yield prediction and growth projection for site-prepared loblolly pine plantations in the Carolinas, Georgia, Alabama and Florida*. University of Georgia School of Forest Resources PMRC Res. Pap. 1996-1, Univ. of Georgia, Athens, GA. 59 p.
- KOZAK, A., AND J.H.G. SMITH. 1993. Standards for evaluating taper estimating systems. *For. Chron.* 69(4):438-444.
- LEDBETTER, J.R., A.D. SULLIVAN, AND T.G. MATNEY. 1986. *Yield tables for cutover site-prepared loblolly pine plantations in the Gulf Coastal Plain*. Mississippi Agric. and For. Exp. Sta. Tech. Bull. 135. 31 p.
- LENHART, J.D. 1996. Total and partial stand-level yield prediction for loblolly and slash pine plantations in east Texas. *South. J. Appl. For.* 20(1):36-41.
- LENHART, J.D., E.V. HUNT, JR., AND J.A. BLACKARD. 1985. Establishment of permanent growth and yield plots in loblolly and slash pine plantations in east Texas. P. 436-437 in *Proc. of the 3rd Biennial Southern Silvicultural Research Conf., 1985*; Shoulders, E. (ed.). US For. Serv. Gen. Tech. Rep. SO-54.
- LENHART, J.D., T.L. HACKETT, C.J. LAMAN, T.J. WISWELL, AND J.A. BLACKARD. 1987. Tree content and taper functions for loblolly and slash pine trees planted on non-old field in east Texas. *South. J. Appl. For.* 10(2):109-112.
- MACKINNEY, A.L., F.X. SCHUMACHER, AND L.E. CHAIKEN. 1937. Construction of yield tables for nonnormal loblolly pine stands. *J. Agric. Res.* 54:531-545.
- MACKINNEY, A.L., AND L.E. CHAIKEN. 1939. *Volume, yield, and growth of loblolly pine in the mid-Atlantic coastal region*. US For. Serv., Appalachian For. Exp. Stn. Tech. Note 33. 30 p.
- MARTIN, S.W., R.L. BAILEY, AND E. JOKELA. 1999. Growth and yield predictions for lower coastal plain slash pine plantations fertilized at mid-rotation. *South. J. Appl. For.* 23(1):39-45.



- OCHI, N., AND Q.V. CAO. 2003. A comparison of compatible and annual growth models. *For. Sci.* 49(2):285–290.
- PIENAAR, L.V., AND W.M. HARRISON. 1989. Simultaneous growth and yield prediction equations for *Pinus elliottii* plantations in Zululand. *S. Afr. For. J.* 149:48–53.
- ROBINSON, A. 2004. Preserving correlation while modelling diameter distributions. *Can. J. For. Res.* 34(1):221–232.
- SCHUMACHER, F.X. 1939. A new growth curve and its application to timber yield studies. *J. For.* 37:819–820.
- SAS INSTITUTE, INC. 2004. *SAS/ETS 9.1 user's guide*. SAS Institute, Inc., Cary, NC. 2,416 p.
- SULLIVAN, A.D., AND J.L. CLUTTER. 1972. A simultaneous growth and yield model for loblolly pine. *For. Sci.* 18(1):76–86.
- ZHAO, D., B. BORDERS, M. WANG, AND M. KANE. 2007. Modeling mortality of second-rotation loblolly pine plantations in the Piedmont/Upper Coastal Plain and Lower Coastal Plain of the southern United States. *For. Ecol. Manag.* 252(1–3):132–143.