

Stephen F. Austin State University

SFA ScholarWorks

Faculty Publications

Chemistry and Biochemistry

2013

A DFT Study of Vibrational Spectra and Mutagenicity Predictions of Mononitrated Fluoranthenes [Abstract]

Kefa Karimu Onchoke

Stephen F Austin State University, onchokekk@sfasu.edu

Jorge Ojeda

Stephen F. Austin State University

Follow this and additional works at: https://scholarworks.sfasu.edu/chemistry_facultypubs



Part of the [Biochemistry Commons](#), and the [Chemistry Commons](#)

[Tell us](#) how this article helped you.

Recommended Citation

Onchoke, Kefa K., and Jorge J. Ojeda. "A DFT Study of Vibrational Spectra and Mutagenicity Predictions of Mononitrated Fluoranthenes." *Polycyclic Aromatic Compounds* 33.5 (2013): 473-500.

This Article is brought to you for free and open access by the Chemistry and Biochemistry at SFA ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SFA ScholarWorks. For more information, please contact cdsscholarworks@sfasu.edu.

A DFT Study of Vibrational Spectra and Mutagenicity Predictions of Mononitrated Fluoranthenes
[Abstract]

Predictions of Mononitrated Fluoranthenes The Raman and infrared (IR) spectra of the mononitrated fluoranthenes (1-, 2-, 3-, 7-, and 8-nitrofluoranthene), important environmental pollutants that induce mutagenic/carcinogenic effects upon mammalian and bacterial cells, were studied via density functional theory, using optimized geometries at the B3LYP/6-311+G(d,p) and PBE0/6-311+G(d,p) levels of theory. The predicted harmonic vibrational frequencies were used in normal mode assignments. To validate the spectra, experimental fluoranthene and 3-nitrofluoranthene were used to benchmark the predicted spectra. The scaled harmonic frequencies below 2000 cm^{-1} are in good agreement to experiment to $<5\text{ cm}^{-1}$ and 10 cm^{-1} or better at the B3LYP/6-311+G(d,p) and PBE0/6-311+G(d,p) levels, respectively. The high intensity nitro asymmetric/symmetric stretch modes ($\nu_{\text{asym}}/\nu_{\text{sym}}\text{NO}_2$), $\nu(\text{C}-\text{C})$ and $\nu(\text{C}-\text{H})$ stretch frequencies were then considered, and used in distinguishing nitrofluoranthene isomers. The $\nu_{\text{sym}}(\text{NO}_2)$ frequency shifts are correlated to the observed mutagenic potencies, thus shedding light on their relative reactivity. The observed differential mutagenic-vibrational correlations, and implications of the ν_{NO_2} shifts/biological mutagenic potencies between nitrofluoranthene isomers are then discussed.