

Stephen F. Austin State University

SFA ScholarWorks

Faculty Publications

Chemistry and Biochemistry

7-27-2011

Experimental and Theoretical Study of Vibrational Spectra of 3-Nitrofluoranthene [Abstract]

Kefa Karimu Onchoke

Stephen F Austin State University, onchokekk@sfasu.edu

Matthew Parks

Stephen F Austin State University

Follow this and additional works at: https://scholarworks.sfasu.edu/chemistry_facultypubs

 Part of the [Chemistry Commons](#)

[Tell us](#) how this article helped you.

Repository Citation

Onchoke, Kefa Karimu and Parks, Matthew, "Experimental and Theoretical Study of Vibrational Spectra of 3-Nitrofluoranthene [Abstract]" (2011). *Faculty Publications*. 42.

https://scholarworks.sfasu.edu/chemistry_facultypubs/42

This Article is brought to you for free and open access by the Chemistry and Biochemistry at SFA ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SFA ScholarWorks. For more information, please contact cdsscholarworks@sfasu.edu.

Experimental and Theoretical Study of Vibrational Spectra of 3-Nitrofluoranthene [Abstract]

The comparative studies of the infrared and Raman spectra of fluoranthene and 3-nitrofluoranthene (3-NF) were made via both FT-IR and density functional theory calculation with a triple ζ Gaussian basis set (B3LYP/6-311 + G(d,p)). The scaled theoretical vibrational frequencies show good agreement to experiment to within $\sim 5 \text{ cm}^{-1}$ for frequencies $< 1800 \text{ cm}^{-1}$. In addition, predictions of the DFT frequencies below 1800 cm^{-1} yielded an overall root mean square (RMS) of ± 4.3 and $\pm 27.4 \text{ cm}^{-1}$ for fluoranthene and 3-NF, respectively. The vibrational assignments are proposed, with particular emphasis on the nitro group vibrations. This study shows the influence the geometrical distortions of the fluoranthene structure have upon nitro group substitution and provides vibrational spectral data useful for the development of spectroscopy-mutagenicity relationships in nitrated polycyclic aromatic hydrocarbons.