

Stephen F. Austin State University

SFA ScholarWorks

---

Faculty Publications

Chemistry and Biochemistry

---

1-2011

## DFT/TD-DFT Investigation of Optical Absorption Spectra, Electron Affinities, and Ionization Potentials of Mono-Nitrated Benzantrones [Abstract]

Kefa Karimu Onchoke

Stephen F Austin State University, onchokekk@sfasu.edu

Follow this and additional works at: [https://scholarworks.sfasu.edu/chemistry\\_facultypubs](https://scholarworks.sfasu.edu/chemistry_facultypubs)

 Part of the [Chemistry Commons](#)

[Tell us](#) how this article helped you.

---

### Repository Citation

Onchoke, Kefa Karimu, "DFT/TD-DFT Investigation of Optical Absorption Spectra, Electron Affinities, and Ionization Potentials of Mono-Nitrated Benzantrones [Abstract]" (2011). *Faculty Publications*. 43.  
[https://scholarworks.sfasu.edu/chemistry\\_facultypubs/43](https://scholarworks.sfasu.edu/chemistry_facultypubs/43)

This Article is brought to you for free and open access by the Chemistry and Biochemistry at SFA ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SFA ScholarWorks. For more information, please contact [cdsscholarworks@sfasu.edu](mailto:cdsscholarworks@sfasu.edu).

## DFT/TD-DFT investigation of optical absorption spectra, electron affinities, and ionization potentials of mono-nitrated benzantrones [Abstract]

Density functional theory (DFT) method B3LYP with basis sets 6-311+G(d,p), 6-31G(d) and 6-311G(d,p) have been used to compute molecular structures, UV-vis absorption spectra, electron affinities (EAs), and ionization potentials (IPs) of benzantrone and 1-, 2-, 3-, 9-, 10-, and 11-nitrobenzantrones. With the use of the time-dependent TD-B3LYP/6-311G(d,p) approach, the UV/vis absorption spectral bands, vertical excitation energies and oscillator strengths was studied. The calculated vertical excitation energies lie within <0.15 eV of the observed values. In comparison to benzantrone, calculated absorption spectra of nitrated nitrobenzantrones show red-shifting to >383 nm, which is correlated to their relative reactivity. Detailed clarification of the singlet-singlet excited states (and energies) are made. The calculated electron affinities are ~0.5 eV greater than in benzantrone (BA); being in the order 3-NBA > 10-NBA > 1-NBA > 2-NBA > 9-NBA > 11-NBA > BA. Similarly, ionization potentials follow a very close order in nitro-BAs. The calculated EAs and IPs implicate both reductive and oxidative pathways on transformation to mutagenic metabolites. The implications of the results are discussed in the context of their differential mutagenic potencies.