

2008

Development and Characterization of Microsatellite Markers for the Grain Amaranths [Abstract]

Melanie A. Mallory

Rozaura V. Hall

Andrea R. McNabb

Donald B. Pratt

Stephen F Austin State University, prattdb@sfasu.edu

Eric N. Jellen

See next page for additional authors

Follow this and additional works at: <https://scholarworks.sfasu.edu/biology>

 Part of the [Biology Commons](#), [Genetics and Genomics Commons](#), and the [Plant Biology Commons](#)

[Tell us](#) how this article helped you.

Repository Citation

Mallory, Melanie A.; Hall, Rozaura V.; McNabb, Andrea R.; Pratt, Donald B.; Jellen, Eric N.; and Maughan, Peter J., "Development and Characterization of Microsatellite Markers for the Grain Amaranths [Abstract]" (2008). *Faculty Publications*. 58.
<https://scholarworks.sfasu.edu/biology/58>

This Abstract is brought to you for free and open access by the Biology at SFA ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SFA ScholarWorks. For more information, please contact cdsscholarworks@sfasu.edu.

Authors

Melanie A. Mallory, Rozaura V. Hall, Andrea R. McNabb, Donald B. Pratt, Eric N. Jellen, and Peter J. Maughan

Development and Characterization of Microsatellite Markers for the Grain Amaranths [Abstract]

The grain amaranths (*Amaranthus hypochondriacus* L., *A. cruentus* L., and *A. caudatus* L.) are important pseudo-cereals native to the Americas. The objective of this project was to produce and characterize a set of highly informative, reproducible microsatellite markers for the grain amaranths. A total of 1457 clones were sequenced from three microsatellite-enriched libraries. Of these, 353 contained unique microsatellites. An additional 29 microsatellite loci were identified from 728 bacterial artificial chromosome–end sequences. A total of 179 microsatellites were polymorphic across accessions from the three grain amaranths. Among these polymorphic microsatellite loci, a total of 731 alleles were identified with an average of four alleles per locus. Heterozygosity values ranged from 0.14 to 0.83, with a mean value of 0.62. Thirty-seven (21%) of the markers were polymorphic between the parents of a segregating population. Phylogenetic analysis using the marker data placed *A. hybridus* L. accessions into two of the three grain amaranth clades, suggesting the polyphyletic evolution of the three cultivated species from different *A. hybridus* ancestors. The transferability of these markers to *A. hybridus*, *A. powellii* S. Wats., and *A. retroflexus* L. is reported and suggests that these markers may be useful in studying other species within the genus *Amaranthus*, including several economically important weeds and ornamentals.