

5-2017

# Observations on Convexity

Chad A. Huckaby

Stephen F Austin State University, chad\_huckaby2000@yahoo.com

Follow this and additional works at: <http://scholarworks.sfasu.edu/etds>



Part of the [Analysis Commons](#)

Tell us how this article helped you.

---

## Repository Citation

Huckaby, Chad A., "Observations on Convexity" (2017). *Electronic Theses and Dissertations*. 88.  
<http://scholarworks.sfasu.edu/etds/88>

This Thesis is brought to you for free and open access by SFA ScholarWorks. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of SFA ScholarWorks. For more information, please contact [cdsscholarworks@sfasu.edu](mailto:cdsscholarworks@sfasu.edu).

---

# Observations on Convexity

## **Creative Commons License**



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

OBSERVATIONS ON CONVEXITY

by

CHAD ALAN HUCKABY, B.S.

Presented to the Faculty of the Graduate School of

Stephen F. Austin State University

In Partial Fulfillment

of the Requirements

For the Degree of

Master of Science

STEPHEN F. AUSTIN STATE UNIVERSITY

May 2017

OBSERVATIONS ON CONVEXITY

by

CHAD ALAN HUCKABY, B.S.

APPROVED:

---

Roy Joseph Harris, Ph.D., Thesis Co-Director

---

Sarah Triana Stovall, Ph.D., Thesis Co-Director

---

Clint Richardson, Ph.D., Committee Member

---

Lee W. Payne, Ph.D., Committee Member

---

Dr. Richard A. Berry, D.M.A.,  
Dean of the Graduate School

## ABSTRACT

This thesis will explore convexity as it pertains to sets of complex-valued functions. These include preliminary looks at established linear and polynomially convex hulls, along with the development of new types of convex hulls. These types will include, but are not limited to the hulls determined by inversions, shift inversions, and Möbius transformations. A convex hull must be preceded by the set of functions involved. These hulls are the smallest convex sets that contain the original set. Justifications and precise definitions are included within the body of the work.

## ACKNOWLEDGEMENTS

There are so many individuals that have played a role in my ability to finish a thesis and a master's degree. As I write this, I hope my acknowledgements are not longer than the actual text of the thesis!

First, I dedicate the entirety of this work to Papa (Marion Huckaby), who passed away just as I was beginning this journey. He was always my biggest fan. He led me to Christ and served in ministry for over 60 years. One day, I hope to write his story, which helps mine be possible.

To my parents, Tim and Monaca: Thank you for teaching me to be humble. My work ethic is due to your influence. No matter the accomplishment, my life has always been focused not on the achievement just received, but the next goal to accomplish even greater things. Thank you for showing me to appreciate the small things in life and not taking things for granted.

To my grandparents, Grandmommy, Nana, and Granddad (Beth, Arlene, and Benny): Thank you for your faith in Christ and showing me that there was no other way. Thank you for your love and support, through whatever means.

To my wife, Candra: Thank you for putting up with my incredibly busy schedule for all of my years at SFA. Your love and compassion has been calming as deadlines for various things have loomed. Thank you for supporting our family as I went to graduate school. You are an amazing teacher and you will be an even better administrator and mother. You are my best friend and I love you with all of my heart. I cannot wait to see what the next chapter of life holds for our family.

To my in-laws, Dee and Rebecca: Thank you first of all for Candra, but also thank you for loving me as your own son. Thank you for supporting me in as many ways

as you were able.

To Steve DaSilva: Thank you for always having the hardest possible science courses at the high school level. You forced your students to seek excellence in a time where standards were continually lowered. I study well because of you. Also, thank you for randomly assigning Candra into my physics lab group right after she moved.

To Dawn Ryan: Thank you for making mathematics fun. You are one of the reasons I chose to study mathematics at SFA. The proofs are for you!

To Lance Taylor: Thank you for helping me to become a better person through athletics. Thank you for fixing my jump shot and letting me pitch. I will never forget my game winning shot against Douglass or my complete game against Timpson.

To Martinsville ISD: Remind your students daily of what is possible with hard work. It does not matter where you attend school. It only matters that you have others that care about your success and the willingness to learn.

To Martinsville Baptist Church: Thank you for allowing me to serve the Lord and His people in our community. I have had incredible experiences due to your support and prayers.

To Scott Sadler: Thanks for being the most supportive brother I could ever hope to have. You have always been the first to recognize my accomplishments and your pride in me has helped my own confidence immensely.

To Casey Duncan: Thanks for being the best (only) male cousin that I have. You are as cool as the other side of the pillow. We both chose the possible wives and they are exactly the same age!

To Joey Bradshaw: Thanks for being my best friend through some of the roughest years of our lives. I am so honored to have served with you at church and in O $\Delta$ K. I am also so glad that you came back to SFA!

To Alli Grimes: Thanks for being there for me in all of our activities over our lives. Thank you for joining O $\Delta$ K and having the impact on our Circle that you did.

Now it is time for you to finish your thesis!

To Brooke Busbee: Thank you for sharing an office with me for the past two years. I am thankful to have gotten to share this experience with you. You will always be my graduate school sister. Your passion for Christ and people is an inspiration to me and to all those around you.

To my mathematics graduate student friends: (Lorna, Greg, Marissa): Thank you for going through this journey with me. I know the work has been hard, the deadlines worse, but your friendship has made the bad times tolerable and the good times great!

To Jamie Bouldin: Thank you for being a great organization adviser, role model, and friend. Thank you for leaving Student Engagement, because you made me grow as a leader by doing so. Thank you for the long conversations about my busy schedule. Thank you for caring so much about the students at SFA and your commitment to excellence in leadership. Thank you for interviewing me for OΔK right after your workout, while drenched in sweat. It may have embarrassed you, but it made me feel at ease, one of your many great qualities.

To Dr. Hollie Smith: Thank you for advising OΔK during Fall 2014. I know how busy you are and you made our transition much easier. Thanks you for your SFA pride and helping select me as Mr. SFA.

To Molly Moody: Thank you for coming in to a new role at SFA and making an immediate impact on student engagement. You challenged me as a leader in almost every aspect, and I have grown so much through that process. You made my tenure as Circle President one of the most enjoyable experiences of my life.

To Dr. Adam Peck: Thank you for being a mentor and friend. I would not be the student leader that I am today without you. Your guidance helped me reach for things that I would previously not have dreamed of. Thank you for caring for the students of SFA in a very real, individual way.



To the rest of the Student Engagement staff: Thank you for the number of hours you put in to make sure that the co-curricular experience at SFA is second to none. You ROCK!

To Dr. Darrell Fry: Thank you for supporting my idea to do chemical demonstrations as research work. Your ability to get the best out of your students is incredible.

To Dr. Michael Tkacik: Thank you for having the hardest undergraduate courses that I ever took. Thank you for helping me succeed at SFA. Thank you for being a mentor, friend, and sounding board for my ideas, both good and bad. You are without a doubt one of the greatest academic influences I have had.

To Dr. Ken Collier: Thank you for teaching me some of the most interesting things I ever learned about politics. Thank you for convincing me that I needed to give more to my studies. Thank you for giving me your book. From that moment, I realized how you viewed me as a student. That gave me a great deal of confidence in my abilities. It may have been a small gesture in your mind, but had a great impact on me.

To Dr. Lee Payne: Thank you for teaching me the quantitative side of political science. The numbers drew me in more than any other subject. Imagine that! Thank you for serving being a friend and mentor as well as serving on my thesis committee. I will beat you at Angry Birds one day!

To Dr. Clint Richardson: Thank you for agreeing to serve on my thesis committee. You were an obvious choice for my topic and I appreciate your knowledge. I wish that I had been in one of your classes. I am thankful that you allowed me to grade for your linear algebra class. Thank you for being a great listener and helping me in many other situations!

To Dr. Pam Roberson: Thank you for being one of the best teachers that I ever had. I can prove things because of you, and it is obvious without proof that you had a huge impact on my education.

To the Omicron Delta Kappa Society: You have helped shape me as a leader. The society's principles are an important part of my life. I am so thankful to have had the opportunity to serve the Stephen F. Austin State University Circle and the national society as a member of the Student Advisory Board and the National Advisory Council.

To Stephen F. Austin State University: Thank you for being the best university in Texas. I cannot imagine having gone anywhere else for either of my degrees. I dread the day that I no longer get to be part of this campus and community. The university has shaped me in so many different areas of my life, I cannot even begin to list them. My pride for SFA will never wane. It can only increase.

To the Board of Regents: Thank you for welcoming me as student regent. You made me feel at ease from the first time I walked into the room. Our university is in capable hands and I cannot wait to see what the future holds for SFA.

To Mrs. Judy Buckingham: Thank you for making my time on the Board of Regents a wonderful experience. Your knowledge and patience made my entire time enjoyable.

To Dr. Baker Pattillo: Thank you for recommending me to Governor Abbott as student regent. Thank you for making my last year as a student at SFA the very best year that I could have imagined. Thank you for your more than fifty years of service to our university. Your leadership of SFA is unmatched by anyone in higher education. Your love for our university has been a catalyst for our growth and has increased my passion for SFA. As legendary as Stephen F. Austin is as the Father of Texas, so you are legendary as president of the university that bears his name. May SFA continue to grow as the best independent university in our state. As always, "Axe 'Em Jacks!"

To Drs. Steve Bullard, Danny Gallant, Steve Westbrook and Mrs. Jill Still: Thank you for showing me the great work that your divisions do at SFA. Thank you

for welcoming me to the team. I am thankful for being able to learn from you during my time as student regent.

To Drs. Harris and Stovall: You are the primary reason I chose to study mathematics. Your passion for teaching and your care for your students make you the best combination of instructors a student could desire. Were it not for taking dual-credit college algebra and trigonometry from the two of you, I would likely have chosen another field of study. I cannot express fully the gratitude that I have for each of you. You have always made me feel comfortable asking questions without fear of being belittled. You have the ability in either the classroom or one-on-one setting to make a student feel valued and intelligent. The fact that you chose to include me in your own research is humbling to me. I have the utmost respect for you and I thank you for working with me. The fact that you are SFA graduates gives you the ability to know where a student is coming from and immediately helps create a connection. My educational experience at SFA has come full circle and it is only fitting that my first two instructors are the ones that will help me complete my master's degree!

To the rest of my friends and family: Thank you for all of your love and support. I could not have done this without you.

Most of all, I thank God for giving me the opportunity to do all of the things I have done at SFA. I thank Him for sending His Son Jesus Christ to die on the cross of Calvary for the remission of my sins.

“I can do all things through Christ who strengthens me”- Philipians 4:13

I believe all victories should end in the SFA Alma Mater:

“All Hail to SFA”  
Oh future bright 'neath the  
Purple and White  
All hail to SFA.  
'Mid Texas pines we have  
Found peaceful shrines  
Where ev'ry month is May.  
Long live our Alma Mater,  
Honor to thee for aye.  
As years unfold, happy  
Mem'ries we'll hold,  
All hail to SFA.

## CONTENTS

<b>ABSTRACT</b> . . . . .	iii
<b>ACKNOWLEDGEMENTS</b> . . . . .	iv
<b>1 INTRODUCTION</b>	<b>1</b>
1.1 Results from Complex Variables . . . . .	1
1.2 Results from Topology . . . . .	3
<b>2 Linear Convex Hull of a Set</b>	<b>8</b>
2.1 Geometric Linear Convex Hull of Compact a Set . . . . .	8
2.2 Functional Linear Convex Hull of a Compact Set . . . . .	12
2.3 Determining the Relationship Between the Functional and Geometric Linear Convex Hull of a Compact Set . . . . .	13
2.4 Linear Convex Hulls of Non-closed, Bounded Sets . . . . .	16
<b>3 Extensions to Other Functional Hulls</b>	<b>24</b>
3.1 Inversion Convex Hull of a Set . . . . .	24
3.2 Shift-Inversion Convex Hull of a Set . . . . .	26
3.3 Möbius Transformation Hull of a Set . . . . .	32
<b>BIBLIOGRAPHY</b> . . . . .	<b>33</b>
<b>VITA</b> . . . . .	<b>34</b>

## LIST OF FIGURES

1.1	$\mathcal{F}$ -Convex Hull diagram for Example 1.5 . . . . .	2
2.1	Construction for the proof of Lemma 2.4 . . . . .	12
2.2	Construction for the proof of Theorem 2.7 . . . . .	14
2.3	Theorem 2.9 - Case 1: Open Disk . . . . .	17
2.4	Theorem 2.9 - Case 2: Non-closed Disk Part I . . . . .	19
2.5	Theorem 2.9 - Case 2: Non-closed Disk Part II . . . . .	20
3.1	Theorem 3.9 Construction . . . . .	28
3.2	Construction for the proof of Theorem 3.11 . . . . .	29

## 1 INTRODUCTION

A common topic in complex variables is the convex hull of a set. The goal of this thesis is to explore different definitions of the convex hull. The functional definition of a convex hull of a set is as follows: Let  $f \in \mathcal{F}$  where  $\mathcal{F}$  is a set of functions and  $\Omega$  is any compact subset of  $\mathbb{C}$ . Then the  $\mathcal{F}$ -convex hull of a set  $\Omega$ , denoted  $\hat{\Omega}_{\mathcal{F}}$ , is defined by

$$\hat{\Omega}_{\mathcal{F}} = \bigcap_{f \in \mathcal{F}} \{w \in \mathbb{C} : |f(w)| \leq M_f\},$$

where  $M_f = \max_{z \in \Omega} |f(z)|$ . This definition requires further definitions to explain, all of which including modulus and compactness, are presented as the remainder of this chapter.

### 1.1 Results from Complex Variables

The usual first definition of convexity is the type called geometric convexity.

**Definition 1.1.** A set  $G$  is **geometrically convex** if given any two points  $a$  and  $b$  in  $G$  the line segment joining  $a$  and  $b$ ,  $\overline{ab}$ , lies entirely in  $G$ .

This type of convexity may also be referred to as linear convexity.

**Definition 1.2.** The **modulus** of a complex number  $z = x + iy$  is defined as

$$|z| = \sqrt{x^2 + y^2},$$

where  $x, y \in \mathbb{R}$ .

**Definition 1.3.** The  $\mathcal{F}$ -**convex hull** of a set is defined as follows: Let  $\mathcal{F}$  be a set of functions and  $\Omega$  any compact subset of  $\mathbb{C}$ . Then the  $\mathcal{F}$ -convex hull of a set  $\Omega$ , denoted  $\hat{\Omega}_{\mathcal{F}}$ , is defined by

$$\hat{\Omega}_{\mathcal{F}} = \bigcap_{f \in \mathcal{F}} \{w \in \mathbb{C} : |f(w)| \leq M_f\},$$

where  $M_f = \max_{z \in \Omega} |f(z)|$ .

*Remark 1.4.* Notice if  $z \in \Omega$ , then  $|f(z)| \leq M_f$  so that  $\Omega \subseteq \{w \in \mathbb{C} : |f(w)| \leq M_f\}$ .

That is,  $\Omega \subseteq \hat{\Omega}_{\mathcal{F}}$ .




Figure 1.1: Example 1.5 Illustration



**Example 1.5.** Let  $\mathcal{F} = \{z, z^2\}$  and  $\Omega = \{z : |z - 1| \leq 1\}$ . Find  $\hat{\Omega}_{\mathcal{F}}$ .

First, it is necessary to find where each of the functions in  $\mathcal{F}$  achieve their respective maxima. On  $\Omega$ , both  $f(z) = z$  and  $g(z) = z^2$  achieve maxima at  $z = 2$ . Then the  $\mathcal{F}$ -convex hull is the intersection of the sets defined by two functions have modulus less than their maxima. For  $f$ , the set of all  $w \in \mathbb{C}$  that causes  $|f(w)| \leq 2$  is the disk centered at the origin of radius 2. For  $g$ , the set of all  $w \in \mathbb{C}$  that causes  $|f(w)| \leq 4$  is also the disk centered at the origin of radius 2. Thus the intersection of these sets is the disk centered at the origin of radius 2. Therefore, the  $\mathcal{F}$ -convex hull of  $\Omega$  is  $\hat{\Omega}_{\mathcal{F}} = \{z : |z| \leq 2\}$ .

## 1.2 Results from Topology

**Definition 1.6.** Let  $X \neq \emptyset$  be a set. Then a **topology** on  $X$  is a collection of subsets of  $X$ , denoted  $\mathcal{T}$ , obeying the following axioms:

- (a)  $X$  and  $\emptyset$  belong to  $\mathcal{T}$ ,
- (b) the intersection of any two elements of  $\mathcal{T}$  is an element of  $\mathcal{T}$ , and
- (c) the union of any sub-collection of  $\mathcal{T}$  is an element of  $\mathcal{T}$ .

**Definition 1.7.** A **topological space** is a set  $X$  together with a topology  $\mathcal{T}$  on  $X$ .

**Definition 1.8.** A set is **open** if it is a member of  $\mathcal{T}$ .

**Definition 1.9.** A set is **closed** if its complement is a member of  $\mathcal{T}$ .

**Definition 1.10.** Let  $X \neq \emptyset$  be a set and  $d : X \times X \rightarrow [0, \infty)$  be a function. Then  $d$  is a **metric** for  $X$  if for any points  $x, y, z \in X$ , the following are true:

- (a)  $d(x, y) \geq 0$
- (b)  $d(x, y) = 0$  if and only if  $x = y$
- (c)  $d(x, y) = d(y, x)$
- (d)  $d(x, z) \leq d(x, y) + d(y, z)$

Note that for this thesis,  $d(z, w) = |z - w|$ .

**Definition 1.11.** Let  $(X, \mathcal{T})$  be a space. A **base** or **basis** for  $\mathcal{T}$  is a collection  $\mathcal{B}$  of subsets of  $X$  such that:

- (a) each member of  $\mathcal{B}$  is also a member of  $\mathcal{T}$ , and
- (b) if  $U \in \mathcal{T}$  and  $U \neq \emptyset$ , then  $U$  is the union of sets belonging to  $\mathcal{B}$ .

**Definition 1.12.** Let  $X$  be a nonempty set of  $\mathbb{C}$  and  $d$  a metric for  $X$ . The unique topology on  $X$  generated by the set of all open  $r$ -spheres in  $X$ , denoted  $B(x, r)$  for some  $r > 0$  and  $x \in X$ , and having these open  $r$ -spheres as a base is called the  ***$d$ -metric topology for  $X$*** . The  $d$ -metric topology is denoted  $\mathcal{T}(d)$ . The topological space  $(X, \mathcal{T})$  is called a ***metric space*** if and only if there exists a metric  $d$  for  $X$  such that the  $d$ -metric topology  $\mathcal{T}(d)$  on  $X$  is the same as  $\mathcal{T}$ . The notation for a metric space is  $(X, d)$ .

**Definition 1.13.** Let  $(X, \mathcal{T})$  be a topological space and  $A \subset X$ . A point  $z$  in set  $X$  is a **boundary point** of  $A$  if and only if every open set in  $X$  containing  $z$  contains at least one point of each of  $X - A$  and  $A$ . The set of boundary points of  $A$  is called the **boundary** of  $A$ .

**Definition 1.14.** Let  $A$  be a subset of a topological space  $X$ . A point  $x \in X$  is a **limit point** of  $A$  if every neighborhood of  $x$  contains at least one point of  $A$  different from  $x$  itself.

**Theorem 1.15.** *A set  $K$  is **open** if and only if it contains none of its boundary points.*

**Theorem 1.16.** *A set  $K$  is **closed** if and only if  $K$  contains all of its boundary points.*

*Remark 1.17.* Note that a set  $K$  is **non-closed** if and only if  $K$  lacks any of its boundary points.

**Definition 1.18.** A set  $K$  is **bounded** if and only if there exists an open disk that contains all of  $K$ .

**Definition 1.19.** Let  $A, B \neq \emptyset$  be two subsets of the metric space  $(X, d)$ . Then the **distance between  $A$  and  $B$** , denoted  $d(A, B)$ , is the greatest lower bound of the set  $\{d(x, y) : x \in A, y \in B\}$ . If  $A = \{a\}$ , this is written  $d(a, B)$  for  $d(A, B)$ .

**Definition 1.20.** Let  $A$  be a set in  $\mathbb{C}$ . The **closure** of  $A$ , denoted  $\bar{A}$ , is defined as follows:

$$\bar{A} = A \cup \{z : z \text{ is a boundary point of } A\}.$$

The following will be especially useful in subsequent work.

**Theorem 1.21.** *Let  $(X, d)$  be a metric space and  $A \neq \emptyset$  a subset of  $X$ . Then  $x \in \bar{A}$  if and only if  $d(x, A) = 0$ .*

*Proof.* Suppose  $x \in \bar{A}$ . Then, for each  $r > 0$ ,  $B(x, r) \cap A \neq \emptyset$  where  $B(x, r)$  is any open disk of radius  $r$  centered at  $x$ . Therefore, for each  $r > 0$  there exists a point  $a_r \in A$  such that  $d(x, a_r) < r$  and, as a consequence, the greatest lower bound of  $\{d(x, a) : a \in A\}$  is zero. The conclusion is that  $d(x, A) = 0$ . For the converse, assume that  $x \in X$  and  $d(x, A) = 0$ . Now if  $x \in A$ , certainly  $x \in \bar{A}$  by the definition of  $\bar{A}$ . So suppose  $x \in X - A$ . It must be shown that  $x$  is a boundary point of  $A$ . Since  $d(x, A) = 0$  for each  $r > 0$  there exists a point  $a_r \in A$  such that  $d(x, a_r) < r$ . It follows that for each  $r > 0$ ,  $B(x, r) \cap A \neq \emptyset$ , showing  $x \in \bar{A}$ . †

**Corollary 1.22.** *If a subset of  $A$  of a metric space  $(X, d)$  is closed and  $x \notin A$ , then  $d(x, A) > 0$ . That is to say from any point not in the closed set, there is a positive distance that exists between the set and that point.*

*Proof.* This result is an immediate consequence of Theorem 1.21. †

**Definition 1.23.** A set  $K$  is compact if and only if every open cover of  $K$  has a finite sub-cover.

**Theorem 1.24** (*Heine-Borel*). *A set  $K$  is compact if and only if  $K$  is both closed and bounded.*

## 2 Linear Convex Hull of a Set

### 2.1 Geometric Linear Convex Hull of Compact a Set

A common interpretation of the linear convex hull is a geometric one. That is for any two points in the set  $\Omega$ , the line segment containing those two points is also in the hull. For this thesis, the following definition is used.

**Definition 2.1.** The **geometric linear convex hull** of a compact set  $\Omega$  is the set composed of all line segments connecting any two points  $z_1, z_2 \in \Omega$ . The geometric linear convex hull of  $\Omega$  is denoted  $\hat{\Omega}_G$ .

The next several pages will include the development of the relationship between the geometric linear convex hull from Definition 2.1 and the functional linear convex hull from Definition 2.5.

**Theorem 2.2.** *A compact set  $\Omega$  is a subset of its geometric linear convex hull,  $\hat{\Omega}_G$ . That is,  $\Omega \subseteq \hat{\Omega}_G$ .*

*Proof.* Let  $z \in \Omega$ . Then  $z$  is either an interior point of  $\Omega$  or  $z$  is some boundary point of  $\Omega$ .

*Case 1:* Suppose that  $z$  is an interior point of  $\Omega$ . Then by definition, there is an open disk  $B(z, r) \subseteq \Omega$ . Then the closed disk  $A = \overline{B(z, \frac{r}{2})} \subseteq \Omega$ . Choose any diameter of  $A$  with points  $z_{1_d}$  and  $z_{2_d}$ . Note that  $z_{1_d}, z_{2_d} \in \Omega$ . Also,  $z \in \overline{z_{1_d}z_{2_d}}$ . Therefore, by

Definition 2.1,  $z \in \hat{\Omega}_G$  and thus  $\Omega \subseteq \hat{\Omega}_G$ .

*Case 2:* Suppose that  $z$  is a boundary point of  $\Omega$ . Let  $z_1$  be any point in  $\Omega$  such that  $z_1 \neq z$ . Then by Definition 2.1,  $\overline{z z_1} \in \hat{\Omega}_G$ . Therefore  $z \in \hat{\Omega}_G$  and  $\Omega \subseteq \hat{\Omega}_G$ .

Therefore it has been shown in both cases that  $\Omega \subseteq \hat{\Omega}_G$ . †

**Theorem 2.3.** *The geometric linear convex hull of a compact set  $\Omega$  is bounded.*

*Proof.* Let  $\Omega$  be a compact set in  $\mathbb{C}$  with geometric linear convex hull  $\hat{\Omega}_G$ . Let  $r = \max_{z \in \Omega} \{|z|\}$ . This maximum exists due to the maximum modulus theorem and the compactness of  $\Omega$ . To prove the result it needs to be the case that  $z_0 \in B(0, r)$  for all  $z_0 \in \hat{\Omega}_G$ . So let  $z_0 \in \hat{\Omega}_G$ . Then there exists  $z_1, z_2 \in \Omega$  such that  $|z_2| \leq |z_1| \leq r$  and  $z_0 \in \overline{z_1 z_2}$ . Since the line segment  $\overline{z_1 z_2}$  can be written as  $z = t z_1 + (1 - t) z_2$  for  $t \in [0, 1]$ , we can write  $z_0 = t z_1 + (1 - t) z_2$  for some  $t \in [0, 1]$ . Now consider the

following inequalities:

$$\begin{aligned}
|z_0| &= |tz_1 + (1-t)z_2| \\
&\leq |tz_1| + |(1-t)z_2| && \text{by the triangle inequality,} \\
&\leq |tz_1| + |(1-t)z_1| && \text{since } |z_1| \geq |z_2|, \\
&= |t||z_1| + |(1-t)||z_1| \\
&= |z_1|(|t| + |1-t|) && \text{by the distributive property,} \\
&= |z_1|(t + 1 - t) && t, (1-t) > 0, t \in \mathbb{R}, \\
&= |z_1|(1) \\
&= |z_1| \\
&\leq r && \text{since } |z_1| \leq r. \\
&< r + 1
\end{aligned}$$

Thus for any  $z_0 \in \hat{\Omega}_G$ ,  $|z_0| \leq r$ . This means for all  $z \in \hat{\Omega}_G$ ,  $z \in B(0, r + 1)$  and  $\hat{\Omega}_G$  is bounded. †

**Lemma 2.4.** *The geometric linear convex hull,  $\hat{\Omega}_G$ , of a compact set  $\Omega$  is closed.*

*Proof.* Let  $z_0$  be a boundary point of  $\hat{\Omega}_G$ . and suppose for contradiction that  $z_0 \notin \hat{\Omega}_G$ . Then there is a sequence  $\{z_n\} \subseteq \hat{\Omega}_G$  such that  $z_n \rightarrow z_0$ . Associated with each  $z_n$ , there is a pair of points  $z_{1_n}, z_{2_n}$  such that  $z_n \in \overline{z_{1_n}z_{2_n}}$  and  $z_{1_n}, z_{2_n} \in \Omega$  (by the definition of geometric hull). Since  $\Omega$  is compact, there is a convergent subsequence  $\{q_{1_n}\}$  of  $\{z_{1_n}\}$ .



Similarly, let  $\{q_{2_n}\}$  be the corresponding subsequence of  $\{z_{2_n}\}$ . Again there is a convergent subsequence  $\{m_{2_n}\}$  of  $\{q_{2_n}\}$  that converges to say  $m_2$ . This means the associated sequence  $\{m_{1_n}\}$  from  $\{q_{1_n}\}$  converges to say  $m_1$ . Define  $\{w_n\}$  from  $\{z_n\}$  to be the corresponding subsequence of points that converges to  $z_0$ . Let  $r = \inf_{z \in \overline{m_1 m_2}} \{|z_0 - z|\}$  and choose  $d = \frac{r}{3}$ . Consider the disks  $\overline{B(z_0, d)}, \overline{B(m_1, d)}, \overline{B(m_2, d)}$ . Let  $l_1$  and  $l_2$  be the common external tangents to  $\overline{B(m_1, d)}, \overline{B(m_2, d)}$ . Define  $t_{11} = l_1 \cap \overline{B(m_1, d)}, t_{12} = l_1 \cap \overline{B(m_2, d)}, t_{21} = l_2 \cap \overline{B(m_1, d)}$ , and  $t_{22} = l_2 \cap \overline{B(m_2, d)}$ . Let  $G = \overline{B(m_1, d)} \cup \overline{B(m_2, d)} \cup \square t_{11} t_{12} t_{22} t_{21}$ . Notice that the minimum distance from  $\overline{B(z_0, d)}$  to  $G$  is  $2d > d$ . So there is an  $N \in \mathbb{N} > 0$  such that for all  $n \geq N$ ,  $|w_n - z_0| < d$ . This means that for all  $n \geq N$ ,  $r = \inf_{z \in \overline{q_{1_n} q_{2_n}}} \{|w_n - z|\} > d$ , but  $\overline{q_{1_n} q_{2_n}} \rightarrow \overline{m_1 m_2}$ , which is a contradiction. Thus  $z_0 \in \hat{\Omega}_G$ . Therefore all boundary points of  $\overline{\hat{\Omega}_G}$  are in  $\hat{\Omega}_G$  and  $\hat{\Omega}_G$  is closed. †




Figure 2.1: Proof of Lemma 2.4.

## 2.2 Functional Linear Convex Hull of a Compact Set

Another interpretation of the linear convex hull is a functional one, for which the definition is given below.

**Definition 2.5.** The **functional linear convex hull** of a compact set  $\Omega$ , denoted  $\hat{\Omega}_{\mathcal{F}}$  is defined as

$$\hat{\Omega}_{\mathcal{F}} = \bigcap_{a,b \in \mathbb{C}} \{w : |aw + b| \leq M_f\},$$

where  $M_f = \max_{z \in \Omega} \{|az + b|\}$ .

*Remark 2.6.* In the complex plane,  $|az + b| \leq r$  means  $|a(z + \frac{b}{a})| \leq r$  or  $|z + \frac{b}{a}| \leq \frac{r}{|a|}$ , which is a closed disk centered at  $-\frac{b}{a}$  of radius  $\frac{r}{|a|}$ . This means that a useful interpretation of Definition 2.5 is that the functional linear convex hull of a set  $\Omega$  is the intersection of all closed disks containing  $\Omega$ .

### 2.3 Determining the Relationship Between the Functional and Geometric Linear Convex Hull of a Compact Set

**Theorem 2.7.** *Let  $\Omega$  be any compact set in  $\mathbb{C}$ . Then the geometric linear convex hull,  $\hat{\Omega}_G$ , is equivalent to the functional linear convex hull, denoted  $\hat{\Omega}_F$ . That is to say  $\hat{\Omega}_G = \hat{\Omega}_F$ .*

*Proof.* Let  $\Omega$  be a compact set in  $\mathbb{C}$ . Let  $\hat{\Omega}_G$  and  $\hat{\Omega}_F$  be the geometric and functional linear convex hulls of  $\Omega$ , respectively.

To show that  $\hat{\Omega}_F \subseteq \hat{\Omega}_G$ , choose a point  $z_0 \notin \hat{\Omega}_G$ . Then from Corollary 1.22, Theorem 2.3, and Lemma 2.4, there exists some minimum distance  $d_0$  from  $z_0$  to the boundary of  $\hat{\Omega}_G$ . Let  $z_1$  be on the boundary of  $\hat{\Omega}_G$  such that  $|z_0 - z_1| = d_0$ . Now, construct the line segment from  $z_0$  to  $z_1$ . Let  $z_2$  be the midpoint of  $\overline{z_0 z_1}$ . Let  $l$  be the line perpendicular to  $\overline{z_0 z_1}$  and passing through  $z_1$ . It can be shown using Euclidean geometry that  $\hat{\Omega}_G$  is contained in the complement of the half-plane that contains  $z_0$ . Since  $\Omega$  is compact, there exists a greatest distance,  $d$ , across  $\Omega$ . Now consider the line  $l$ , containing  $z_1$  perpendicular to  $\overline{z_0 z_1}$  and choose points  $z_{1_d}, z_{2_d}$  on this line a distance of  $d$  away from

$z_1$  so that  $z_1$  is between  $z_{1_d}$  and  $z_{2_d}$ . Since three distinct noncollinear points uniquely determine a circle, construct the disk determined by the circle formed by  $z_2, z_{1_d}, z_{2_d}$ . This disk captures all of  $\hat{\Omega}_G$  and excludes  $z_0$ . Thus  $z_0$  can be removed from contention for membership in  $\hat{\Omega}_{\mathcal{F}}$ . Since  $z_0$  was arbitrary, any point not in  $\hat{\Omega}_G$  will also not be in  $\hat{\Omega}_{\mathcal{F}}$ .




Figure 2.2: Proof of Theorem 2.7.

To show  $\hat{\Omega}_G \subseteq \hat{\Omega}_{\mathcal{F}}$ , let  $z \in \hat{\Omega}_G$ . Then there are two points  $z_1, z_2 \in \Omega$  such that  $z \in \overline{z_1 z_2}$ . Then the segment  $\overline{z_1 z_2}$  can be written as

$$z = tz_1 + (1 - t)z_2,$$

for  $t \in [0, 1]$ . Let  $B(q, r)$  be any disk containing  $\Omega$ . Without loss of generality, let

$|z_2 - q| \leq |z_1 - q| \leq r$ . Now consider the following inequalities for any  $z \in \overline{z_1 z_2}$ .

$$\begin{aligned}
|z - q| &= |t(z_1 - q) + (1 - t)(z_2 - q)| \\
&\leq |t(z_1 - q)| + |(1 - t)(z_2 - q)| && \text{by the triangle inequality,} \\
&\leq |t(z_1 - q)| + |(1 - t)(z_1 - q)| && \text{since } |(z_1 - q)| \geq |(z_2 - q)|, \\
&= |t|(z_1 - q)| + |(1 - t)|(z_1 - q)| \\
&= |(z_1 - q)|(|t| + |1 - t|) && \text{by the distributive property,} \\
&= |(z_1 - q)|(t + 1 - t) && t, (1 - t) > 0, t \in \mathbb{R}, \\
&= |(z_1 - q)|(1) \\
&= |(z_1 - q)| \\
&\leq r && \text{since } |(z_1 - q)| \leq r.
\end{aligned}$$

Thus every point  $z \in \overline{z_1 z_2}$  is contained in *every* closed disk containing  $\Omega$ . Therefore  $z_0 \in \hat{\Omega}_{\mathcal{F}}$ . Therefore, both containments have been shown and for compact sets  $\Omega$  in  $\mathbb{C}$ ,

$$\hat{\Omega}_G = \hat{\Omega}_{\mathcal{F}}.$$

†

Why is there interest in looking at the relationship between geometric and functional linear convex hulls? As shown in the previous result, compactness makes the

respective hulls equal. Relaxing the condition of compactness leads to some interesting results.

## 2.4 Linear Convex Hulls of Non-closed, Bounded Sets

**Definition 2.8.** The **geometric linear convex hull** of *any* set  $\Omega$  is the set composed of all line segments connecting any two points  $z_1, z_2 \in \Omega$ . The geometric linear convex hull of  $\Omega$  is denoted  $\hat{\Omega}_G$ .

To show that  $\hat{\Omega}_G$  is not necessarily equal to  $\hat{\Omega}_F$  for any set in  $\mathbb{C}$ , consider the following example.

**Theorem 2.9.** *Let  $\Omega$  be a subset of the closed unit disk centered at the origin containing all of the interior of the disk. Then one of the following is true:*

1. *If  $\Omega$  is open, then the geometric linear convex hull is itself. That is  $\Omega = \hat{\Omega}_G$*
2. *If  $\Omega$  contains any number of its boundary points, where  $\mathbf{B}$  is some subset of the boundary of  $\Omega$ , then the geometric linear convex hull will be the interior of the disk with the boundary points added. That is  $\hat{\Omega}_G = \Omega \cup \mathbf{B}$ .*




Figure 2.3: Case 1.

*Case 1: Let  $\Omega$  be open. Then the geometric linear convex hull of  $\Omega$  is itself.*

*Proof.* Let  $\Omega$  be open. Further let  $z_1, z_2 \in B(0, 1)$ . Then any point  $z$  on the line segment  $\overline{z_1 z_2}$  can be written in the following way:

$$z = z_1 \cdot (t) + (1 - t) \cdot z_2 \text{ for some } t \in [0, 1]$$

That is  $\overline{z_1 z_2} = \{z : z = z_1 \cdot (t) + (1 - t) \cdot z_2 \text{ for } t \in [0, 1]\}$ . Here, it is necessary to show that the entire segment is contained within  $\Omega$ . That is to say, it is necessary to show the following inequality:

$$|z_1 \cdot (t) + (1 - t) \cdot z_2| < 1$$

Now assume without loss of generality that  $|z_1| \geq |z_2|$ . Consider the following:

$$\begin{aligned}
|z_1 \cdot t + (1-t) \cdot z_2| &\leq |t \cdot z_1| + |(1-t) \cdot z_2| && \text{by the triangle inequality,} \\
&= |t| \cdot |z_1| + |(1-t)| \cdot |z_2| && \text{since } |ab| = |a| \cdot |b|, \\
&\leq |t| \cdot |z_1| + |(1-t)| \cdot |z_1| && \text{by assumption that } |z_1| > |z_2|, \\
&= |z_1| (|t| + |1-t|) \\
&= |z_1| (t + 1 - t) && \text{since } t, 1-t \text{ are positive,} \\
&= |z_1| \cdot (1) \\
&= |z_1| \\
&< 1 && \text{since } |z_1| < 1.
\end{aligned}$$

Thus every point on every line segment has modulus strictly less than 1. That is to say that every point is on the interior of  $B(0, 1)$ . Therefore the geometric linear convex hull of the open unit disk is itself.

†

*Case 2: Let  $\Omega$  contain any number of its boundary points. Let  $z_1$  and  $z_2$  be in  $\Omega$ . If  $z_1$  and  $z_2$  are interior points, the previous proof shows that  $\overline{z_1 z_2} \subseteq \Omega$ . If not, either  $z_1, z_2$  are both boundary points or one is a boundary point and the other is an interior point.*






Figure 2.4: Non-closed disk Part I: Case 2

*Proof.* First, consider that  $z_1$  is a boundary point and  $z_2$  is an interior point of  $\Omega$ . Then by assumption  $|z_1| = 1$  and  $|z_2| < 1$ . Then any point  $z$  on the line segment  $\overline{z_1 z_2}$  can be written in the following way:

$$z = z_1 \cdot (t) + (1 - t) \cdot z_2 \text{ for some } t \in [0, 1].$$

Here, it is necessary to show that the only boundary points that are picked up in the hull for  $\Omega$  are the boundary points in  $\Omega$ . It is necessary to show the inequality below, with equality only at  $t = 1$ .

$$|z_1 \cdot (t) + (1 - t) \cdot z_2| \leq 1$$

Consider the following:

$$\begin{aligned}
 |z_1 \cdot t + (1-t) \cdot z_2| &\leq |t \cdot z_1| + |(1-t) \cdot z_2| && \text{by the triangle inequality,} \\
 &= |t| \cdot |z_1| + |(1-t)| \cdot |z_2| && \text{since } |ab| = |a| \cdot |b|, \\
 &< |t| \cdot |z_1| + |(1-t)| \cdot |z_1| && \text{by assumption that } |z_1| > |z_2|, \\
 &= (|t| + |1-t|) && \text{since } |z_1| = 1 \\
 &= (t + 1 - t) && \text{since } t, 1-t \text{ are positive,} \\
 &= 1
 \end{aligned}$$

Thus every point on every line segment has modulus less than or equal to 1. Equality occurs at  $t = 1$ , which is the boundary point  $z_1$ .




Figure 2.5: Non-closed disk Part II: Case 2

Now suppose  $z_1, z_2$  are on the boundary of  $\Omega$ . By assumption  $|z_1| = 1$  and  $|z_2| = 1$ .

Then any point  $z$  on the line segment  $\overline{z_1 z_2}$  can be written in the following way:

$$z = z_1 \cdot (t) + (1 - t) \cdot z_2 \text{ for some } t \in [0, 1].$$

Here, it is necessary to show that the segment  $\overline{z_1 z_2}$  contains only interior points other than the endpoints. That is, it must be shown that

$$|z_1 \cdot (t) + (1 - t) \cdot z_2| \leq 1, \text{ with equality only occurring at the endpoints.}$$

Consider the following:

$$\begin{aligned} |z_1 \cdot (t) + (1 - t) \cdot z_2| &\leq |t \cdot z_1| + |(1 - t) \cdot z_2| && \text{by the triangle inequality,} \\ &= |t| \cdot |z_1| + |(1 - t)| \cdot |z_2| && \text{since } |ab| = |a| \cdot |b|, \\ &= |t| + |(1 - t)| && \text{by assumption that } |z_1| = |z_2| = 1, \\ &= (|t| + |1 - t|) \\ &= (t + 1 - t) && \text{since } t, 1 - t \text{ are positive,} \\ &= 1 \end{aligned}$$

Thus every point on every line segment has modulus less than or equal to 1. Equality occurs at  $t = 1$  (the endpoint  $z_1$ ) and at  $t = 0$  (the endpoint  $z_2$ ). †

It has been shown that for any non-closed subset of the unit disk containing the interior of the disk and centered at the origin, that the geometric linear convex hull is the original set itself.

**Theorem 2.10.** *For the functional case, let  $\Omega = B(0,1) \cup \mathbf{B}$ , where  $\mathbf{B}$  is some subset of the boundary of  $B(0,1)$ . Then the functional linear convex hull of  $\Omega$  is the closed unit disk,  $\bar{\Omega}$ . That is  $\hat{\Omega}_{\mathcal{F}} = \bar{\Omega}$ .*

*Proof.* Let  $\Omega = B(0,1) \cup \mathbf{B}$ . Then the functional linear convex hull of  $\Omega$  is the intersection of all disks containing all of  $\Omega$ . To show  $\hat{\Omega}_{\mathcal{F}} \subseteq \bar{\Omega}$ , let  $z \in \hat{\Omega}_{\mathcal{F}}$ . Then  $z$  is in the intersection of all closed disks containing  $\Omega$ . Since  $\bar{\Omega}$  is a closed disk containing  $\Omega$ ,  $z \in \bar{\Omega}$ . Thus  $\hat{\Omega}_{\mathcal{F}} \subseteq \bar{\Omega}$ . To show  $\bar{\Omega} \subseteq \hat{\Omega}_{\mathcal{F}}$ , let  $z \in \bar{\Omega}$ . Then either  $z \in \Omega$  or  $z \notin \Omega$  but  $z$  is a boundary point of  $\Omega$ .

*Case 1:* If  $z \in \Omega$ , then  $z \in \Omega \subseteq \hat{\Omega}_{\mathcal{F}}$ . Thus  $z \in \hat{\Omega}_{\mathcal{F}}$ .

*Case 2:* If  $z$  is a boundary point of  $\Omega$  and  $z \notin \Omega$ , then  $z \in \overline{B(0,1)}$ . Note that the closed unit disk centered at the origin is in the intersection of all of the closed disks containing  $\Omega$ . Thus,  $z \in \hat{\Omega}_{\mathcal{F}}$ . In either case,  $\bar{\Omega} \subseteq \hat{\Omega}_{\mathcal{F}}$ .

Therefore  $\hat{\Omega}_{\mathcal{F}} = \bar{\Omega}$ . †

To summarize, the functional linear convex hull is not always the same as the geometric linear convex hull for non-closed, bounded sets. This is the case because of the definition of the functional linear convex hull. The functional linear hull is defined as the intersection of all of the disks containing the original set. An arbitrary intersection of closed sets is closed, which implies that the functional hull is closed. This is in contrast to the geometric linear convex hull of the set in Theorem 2.9 and Theorem 2.10. The functional and geometric linear convex hulls always agree when

the original set is compact.

### 3 Extensions to Other Functional Hulls

In the previous chapters, finding the  $\mathcal{F}$ -hull of a set required a set of functions  $\mathcal{F}$  and a compact set  $\Omega \subseteq \mathbb{C}$ . Note that the functions that comprised  $\mathcal{F}$  have been entire. For the remaining sections, the assumption is that all of the functions in  $\mathcal{F}$  are analytic on  $\Omega$ .

#### 3.1 Inversion Convex Hull of a Set

The inversion convex hull of a set is found using the functions that consist of inversions of a complex numbers of the form  $f(z) = \frac{a}{z}$ , where  $a \in \mathbb{C}$  is a constant and  $z \in \mathbb{C}$  is a variable. It is useful to begin the exploration with the definition of the inversion convex hull of a set.

**Definition 3.1.** Let  $\Omega \subseteq \mathbb{C} - \{0\}$  be compact and  $\mathcal{I} = \left\{ f(z) : f(z) = \frac{a}{z}; a \in \mathbb{C} \right\}$ .

The **inversion convex hull** of a set  $\Omega$ , denoted  $\hat{\Omega}_{\mathcal{I}}$ , is defined as:

$$\hat{\Omega}_{\mathcal{I}} = \bigcap_{f \in \mathcal{I}} \{w \in \mathbb{C} : |f(w)| \leq M_f\},$$

where  $M_f = \max_{z \in \Omega} \{|f(z)|\}$ .

Now that there is some basis to find an inversion convex hull of a set, begin with a finite set and find its hull.

**Example 3.2.** Consider the set  $\Omega = \{1, 2\}$ . Then for any  $a \in \mathbb{C}$ ,  $\max_{z \in \Omega} \left\{ \left| \frac{a}{z} \right| \right\} = |a|$ . That is for  $|z| \geq 1$ ,  $\left| \frac{a}{z} \right| \leq \max_{z \in \Omega} \left\{ \left| \frac{a}{z} \right| \right\} = |a|$  and for  $|z| < 1$ ,  $\left| \frac{a}{z} \right| > |a|$  when  $a \neq 0$ . Therefore the inversion convex hull for  $\Omega$  is  $\hat{\Omega}_{\mathcal{I}} = \{|z| \geq 1; z \in \mathbb{C}\}$ . A more visual description of this set is the complex plane with the open unit disk removed.

The previous example gives some intuition as to what the inversion convex hull of any two point set may be.

**Theorem 3.3.** *Let  $\Omega = \{z_1, z_2\}$  be a two point set in  $\mathbb{C} - \{0\}$ . Then the inversion convex hull of the set  $\Omega$  is the set  $\hat{\Omega}_{\mathcal{I}} = \{|z| \geq r\}$  where  $r = \min\{|z_1|, |z_2|\} \leq |z_i|$  for all  $z_i \in \Omega$ .*

*Proof.* Let  $\Omega = \{z_1, z_2\}$ , where  $z_1, z_2 \in \mathbb{C}$  and  $z_1, z_2 \neq 0$ . Without loss of generality, let  $r = |z_1| \leq |z_2|$  and let  $K = \{z : |z| \geq |z_1|\}$ . Let  $z_0 \in K$ . Note that  $\max_{z \in \Omega} \left\{ \left| \frac{a}{z} \right| \right\} = \left| \frac{a}{z_1} \right| = \frac{|a|}{r}$ . Then  $\left| \frac{a}{z_0} \right| \leq \left| \frac{a}{z_1} \right|$  and thus  $z_0 \in \{w \in \mathbb{C} : |f(w)| \leq \left| \frac{a}{z_0} \right|\}$  is in the inversion convex hull of  $\Omega$ . Now let  $w$  be in the inversion convex hull of  $\Omega$ . That is to say  $\left| \frac{a}{w} \right| \leq \max_{z \in \Omega} \left\{ \left| \frac{a}{z} \right| \right\} = \left| \frac{a}{z_1} \right| = |a|$  for all  $a \in \mathbb{C}$ . Therefore  $|w| \geq |z_1|$  and thus  $w \in K$ . Therefore,  $K$  is the inversion convex hull of  $\Omega$ . †

**Theorem 3.4.** *Let  $\Omega = \{z_1, z_2, \dots, z_n\}$  be any finite set in  $\mathbb{C} - \{0\}$ . Then the inversion convex hull of the set  $\Omega$  is the set  $\hat{\Omega}_{\mathcal{I}} = \{|z| \geq r\}$  where  $r = \min\{|z_1|, |z_2|, \dots, |z_n|\} \leq |z_i|$  for all  $z_i \in \Omega$ .*

*Proof.* Let  $\Omega = \{z_1, z_2, \dots, z_n\}$ , where  $z_1, z_2, \dots, z_n \in \mathbb{C}$  and  $z_1, z_2, \dots, z_n \neq 0$ . Without loss of generality, let  $|z_1| \leq |z_i|$  for all  $1 \leq i \leq n$ . Then let  $K = \{z : |z| \geq |z_1|\}$ . Let

$z_0 \in K$ . Note that  $\max_{z \in \Omega} \left\{ \left| \frac{a}{z} \right| \right\} = \left| \frac{a}{z_1} \right| = |a|$  for  $z \in \Omega$ . Then  $\left| \frac{a}{z_0} \right| \leq \left| \frac{a}{z_1} \right|$  and thus  $z_0$  is in the inversion convex hull of  $\Omega$ ,  $\hat{\Omega}_{\mathcal{I}}$ . Conversely, let  $w$  be in the inversion convex hull of  $\Omega$ . That is to say  $\left| \frac{a}{w} \right| \leq \max_{z \in \Omega} \left\{ \left| \frac{a}{z} \right| \right\} = \left| \frac{a}{z_1} \right|$  for all  $w \in \Omega$ . Therefore  $|w| \geq |z_1|$  and thus  $w \in K$ . Therefore,  $K$  is the inversion convex hull of  $\Omega$ . †

**Theorem 3.5.** *Let  $\Omega$  be a closed set in  $\mathbb{C} - \{0\}$ . Then the inversion convex hull of  $\Omega$  is  $\hat{\Omega}_{\mathcal{I}} = \mathbb{C} - B(0, r)$ , where  $r$  is the distance from 0 to the nearest point in  $\Omega$ .*

*Proof.* Using the same argument as the proof of Theorem 3.3, choose whichever point in  $\Omega$  that is closest to the origin. This point is guaranteed to exist by Corollary 1.22. The result follows. †

### 3.2 Shift-Inversion Convex Hull of a Set

The shift-inversion convex hull of a set is found using the functions that consist of inversions of a complex numbers of the form  $f(z) = \frac{a}{z-b}$ , where  $a, b \in \mathbb{C}$  are constants and  $z \in \mathbb{C}$  is a variable. It is useful to begin with the definition of the shift-inversion convex hull of a set.

**Definition 3.6.** Let  $\Omega \subseteq \mathbb{C}$  be compact and  $\mathcal{S} = \left\{ f(z) : f(z) = \frac{a}{z-b}; a, b \in \mathbb{C} \right\}$ . The **shift-inversion convex hull** of a set  $\Omega$ , denoted  $\hat{\Omega}_{\mathcal{S}}$ , is defined as:

$$\hat{\Omega}_{\mathcal{S}} = \bigcap_{f \in \mathcal{S}} \{w \in \mathbb{C} : |f(w)| \leq M_f\},$$

where  $M_f = \max_{z \in \Omega} \{|f(z)|\}$ .



Now to get to the desired shift-inversion convex hull, the more useful way to proceed is to negate Definition 3.6 to find the complement of  $\hat{\Omega}_{\mathcal{S}}$ ,  $\hat{\Omega}_{\mathcal{S}}^c = \mathbb{C} - \hat{\Omega}_{\mathcal{S}}$ .

**Definition 3.7.** The complement of the shift-inversion convex hull of a set  $\Omega$  is the set  $\mathbb{C} - \hat{\Omega}_{\mathcal{S}}$  defined as

$$\hat{\Omega}_{\mathcal{S}}^c = \bigcup_{f \in \mathcal{S}} \{w : |f(w)| > M_f\},$$

where  $M_f = \max_{z \in \Omega} \{|f(z)|\}$ .

The above definition allows the verification that a point  $z_0$  is not in the hull by finding a single shift-inversion  $f$  so that  $|f(z_0)| > \max_{z \in \Omega} \{|f(z)|\}$ . With this knowledge, consider the following example.

**Example 3.8.** Let  $\Omega = \{1, 2\}$ . Then the shift-inversion convex hull of the set  $\Omega$  is the original set  $\Omega = \{1, 2\}$ .

*Proof.* Let  $\Omega = \{1, 2\}$  and let  $z_0 \in \mathbb{C} - \Omega$ . Define  $d_1 = |z_0 - 1|$ ,  $d_2 = |z_0 - 2|$ , and let  $\delta = \min\{d_1, d_2\}$ . Suppose without loss of generality that  $\delta = d_2$ . Choose  $z_\alpha$  on  $\overline{2z_0}$  such that  $|z_0 - 2| > |z_0 - z_\alpha|$ . Choose  $b$  on the ray  $\overrightarrow{z_\alpha z_0}$  such that  $|b - z_0| = \frac{\delta}{4}$  and  $|b - z_\alpha| > |z_\alpha - z_0|$ . Consider  $f(z) = \frac{1}{z - b}$ . It is necessary to show that  $|f(z_0)| > |f(z_\alpha)|$ . Based on the choice of  $b$ , it is the case that  $\frac{1}{|z_0 - b|} > \frac{1}{|z_\alpha - b|}$ . Therefore, the only points that are in the shift-inversion convex hull are points that are in  $\Omega$  to begin with. That is to say that  $\Omega$  is its own shift-inversion convex hull. †




Figure 3.1: Construction for Theorem 3.9.

**Theorem 3.9.** *Let  $\Omega$  be any finite two point set. Then the shift-inversion convex hull of the set  $\Omega$  is the original set  $\Omega$ .*

*Proof.* Let  $\Omega = \{z_1, z_2\}$  and let  $z_0 \in \mathbb{C} - \Omega$ . Then  $d_1 = |z_0 - z_1|$  and  $d_2 = |z_0 - z_2|$ . Let  $\delta = \min\{d_1, d_2\}$ . Suppose without loss of generality that  $\delta = d_2$ . Choose  $z_\alpha \neq z_0$  on  $\overline{z_2 z_0}$  such that  $|z_0 - z_2| > |z_0 - z_\alpha|$ . Choose  $b$  on the ray  $\overrightarrow{z_\alpha z_0}$  such that  $|b - z_0| = \frac{\delta}{4}$  and  $|b - z_\alpha| > |z_\alpha - z_0|$ . Consider  $f(z) = \frac{1}{z - b}$ . It is necessary to show that  $|f(z_0)| > |f(z_\alpha)|$ . Based on the choice of  $b$ , it is the case that  $\frac{1}{|z_0 - b|} > \frac{1}{|z_\alpha - b|}$ . Therefore, the only points that are in the shift-inversion convex hull are points that are in  $\Omega$  to begin with. That is to say that  $\Omega$  is its own shift-inversion convex hull. †

**Theorem 3.10.** *Let  $\Omega$  be any finite set. Then the shift-inversion convex hull of the set  $\Omega$  is the original set,  $\Omega$ .*

*Proof.* Let  $\Omega = \{z_1, z_2, \dots, z_n\}$  where  $z_i \in \mathbb{C}$  for all  $1 \leq i \leq n$  and let  $z_0 \in \mathbb{C} - \Omega$ . Let  $d_i = |z_0 - z_i|$  for all  $1 \leq i \leq n$  and  $\delta = \min\{d_i\}$ . Suppose without loss of generality that  $\delta = d_1$ . Let  $b \in \overline{z_0 z_1}$  such that  $b \neq z_0$  and  $|b - z_0| = \frac{\delta}{4}$ . Then the following is true:

$$\begin{aligned}
 \delta &= |z_0 - b| + |b - z_1| && \text{Since } b \in \overline{z_0 z_1} \\
 &\leq |z_0 - z_i| && \text{Since } \delta \text{ is minimal over } \Omega, \\
 &\leq |z_0 - b| + |b - z_i| && \text{by the triangle inequality,}
 \end{aligned}$$

Also,  $|b - z_1| \leq |b - z_i|$  and since  $b$  is chosen closer to  $z_0$  than  $z_1$ ,  $|z_0 - b| < |b - z_1|$ , so  $|z_0 - b| < |b - z_1| \leq |b - z_i|$ . The first and last parts of the previous inequality show that the point  $z_0$  is closer to  $b$  than any point  $z_i \in \Omega$ . Consider the function  $f(z) = \frac{1}{z - b}$ . Then  $|f(z_0)| = \frac{1}{|z_0 - b|} > \frac{1}{|z_i - b|} = |f(z_i)|$  for all  $z_i \in \Omega$ . Therefore, the only points that are in the shift-inversion convex hull are points that are in  $\Omega$  to begin with. That is to say that  $\Omega$  is its own shift-inversion convex hull. †




Figure 3.2: Constructions for Theorem 3.11

**Theorem 3.11.** *Let  $\Omega$  be any closed bounded simply connected set. Then the shift-inversion convex hull of the set  $\Omega$  is the original set,  $\Omega$ .*

*Proof.* Let  $\Omega$  be any closed bounded simply connected set and let  $z_0 \in \mathbb{C} - \Omega$ . The following constructions can be seen in Figure 3.2. From Theorem 1.21 and Corollary 1.22, there exists a smallest distance from  $z_0$  to a point  $z_1 \in \Omega$ , which is on the boundary of  $\Omega$ . Let this smallest distance be called  $\delta$ . Consider  $B(z_0, \frac{\delta}{4})$  and choose  $b \in \overline{z_0 z_1} \cap B(z_0, \frac{\delta}{4})$ , where  $b \neq z_0$ . Consider  $f(z) = \frac{1}{z - b}$ . It is necessary to show that  $|f(z_0)| > |f(z_1)| \geq |f(z)|$  for all  $z \in \Omega$ . Based on the choice of  $b$ , it is the case that  $\frac{1}{|z_0 - b|} > \frac{1}{|z_1 - b|} \geq \frac{1}{|z - b|}$  for all  $z \in \Omega$ . Therefore, the only points that are in the shift-inversion convex hull are points that are in  $\Omega$  to begin with. That is to say that  $\Omega$  is its own shift-inversion convex hull. †

**Theorem 3.12.** *Any closed set is its own shift inversion convex hull.*

*Proof.* Let  $\Omega$  be a closed set and  $\hat{\Omega}_S$  be its shift-inversion convex hull. Then by definition of shift-inversion convex hull,  $\Omega \subseteq \hat{\Omega}_S$ . Now for the other containment, assume for contradiction that  $z_0 \in \hat{\Omega}_S$  and  $z_0 \notin \Omega$ . As in a previous proof in this thesis, there is a neighborhood around  $z_0$  such that a specific shift-inversion to exclude  $z_0$  from the hull can be found. This contradicts  $z_0 \in \hat{\Omega}_S$ . Therefore all  $z \in \hat{\Omega}_S$  are also in  $\Omega$ , or  $\hat{\Omega}_S \subseteq \Omega$ . Therefore  $\Omega = \hat{\Omega}_S$  and every closed set is its own shift-inversion convex hull. †

**Theorem 3.13.** *The shift-inversion convex hull of an open set is the closure of the set.*

*Proof.* Let  $\Omega$  be an open set and  $\hat{\Omega}_{\mathcal{S}}$  be its shift-inversion convex hull. Let  $z_1$  be a boundary point for  $\Omega$ . Then there is no neighborhood around  $z_1$  where a shift-inversion to exclude  $z_1$  from  $\hat{\Omega}_{\mathcal{S}}$  could be found because any neighborhood would also include points inside  $\Omega$  by Definition 1.13, which cannot be excluded. Thus every boundary point of an open set is included in  $\hat{\Omega}_{\mathcal{S}}$ . Therefore  $\hat{\Omega}_{\mathcal{S}}$  is closed. Since  $\hat{\Omega}_{\mathcal{S}}$  is closed, its shift-inversion convex hull is  $\hat{\Omega}_{\mathcal{S}}$ . Thus the shift-inversion convex hull of  $\Omega$  is  $\hat{\Omega}_{\mathcal{S}} = \bar{\Omega}$ . †

**Corollary 3.14.** *The shift-inversion convex hull of a non-closed set is the closure of the set.*

*Proof.* For boundary points, use the same argument used in proof of Theorem 3.13. For points outside the boundary, use the same argument used in the proof of Theorem 3.12. †

**Corollary 3.15.** *A shift-inversion convex hull is always a closed set.*

*Proof.* The result follows from the fact that closed and non-closed sets have shift-inversion convex hulls that are closed. †

*Remark 3.16.* From the previous work in this section there is a complete classification of the shift-inversion convex hull for *any* set. In summary, for any set  $\Omega \in \mathbb{C}$ ,  $\hat{\Omega}_{\mathcal{S}} = \bar{\Omega}$ .

### 3.3 Möbius Transformation Hull of a Set

The Möbius transformation convex hull of a set is found using the set of functions are called Möbius transformations, that is, functions of the form  $f(z) = \frac{az + b}{cz + d}$ , where  $a, b, c, d \in \mathbb{C}$  are constants and  $z \in \mathbb{C}$  is a variable with  $ad - bc \neq 0$ . Below is the definition of the shift-inversion convex hull of a set.

**Definition 3.17.** Let  $\Omega \subseteq \mathbb{C}$  be compact and  $\mathcal{M} = \left\{ f(z) : f(z) = \frac{az + b}{cz + d}; a, b, c, d \in \mathbb{C} \right\}$ .

The **Möbius transformation hull** of a Set of a set  $\Omega$ , denoted  $\hat{\Omega}_{\mathcal{M}}$ , is defined as:

$$\hat{\Omega}_{\mathcal{M}} = \bigcap_{f \in \mathcal{M}} \{w \in \mathbb{C} : |f(w)| \leq M_f\},$$

where  $M_f = \max_{z \in \Omega} \{|f(z)|\}$ .

**Theorem 3.18.** *A closed set  $\Omega$  is its own Möbius transformation convex hull .*

*Proof.* Given that all shift-inversions are Möbius transformations and that  $\Omega$  is a subset of its  $\mathcal{M}$ -convex hull, then the result is proven. †

The previous result leads to a larger, more general result for  $\mathcal{F}$ -convex hulls.

**Theorem 3.19.** *Suppose two sets of functions have the following relationship:  $\mathcal{F}_1 \subseteq \mathcal{F}_2$ . Then the convex hull of a set  $\Omega$  has the relationship  $\hat{\Omega}_{\mathcal{F}_2} \subseteq \hat{\Omega}_{\mathcal{F}_1}$*

*Proof.* Let  $\mathcal{F}_1 \subseteq \mathcal{F}_2$ . Let  $\Omega$  be any set and  $\hat{\Omega}_{\mathcal{F}_1}$  and  $\hat{\Omega}_{\mathcal{F}_2}$  be the hulls under the respective sets of functions. Since there is potential for more functions in  $\mathcal{F}_2$ , there is a chance to exclude more points from the hull. Therefore  $\hat{\Omega}_{\mathcal{F}_2} \subseteq \hat{\Omega}_{\mathcal{F}_1}$ . †

## BIBLIOGRAPHY

- [1] John B. Conway, *Functions of one complex variable*, Springer-Verlag, 1978.
- [2] Paul E. Long, *An introduction to general topology*, Charles E. Merrill Publishing Company, 1971.

## VITA

Chad Huckaby was born in Nacogdoches, Texas on 20 October 1992. He is a proud native East Texan. He is also a proud third generation SFA Lumberjack. He attended Martinsville High School in Martinsville, Texas, USA, graduating as Valedictorian in 2011. He attended Stephen F. Austin State University in Nacogdoches, Texas from 2011-2015 and received a Bachelor of Science degree in Mathematics and Political Science, with a minor in Chemistry in May 2015. In 2015, He was named Mr. SFA for his dedication and service to Stephen F. Austin State University. He began graduate studies in Mathematical Sciences at Stephen F. Austin State University in Fall 2015 and is expected to graduate in May 2017. In 2016, he was appointed by Texas Governor Greg Abbott to serve the State of Texas as the student representative on the Board of Regents of Stephen F. Austin State University for the academic year 2016-2017. When his studies are complete at SFA, he intends to pursue a doctoral degree in curriculum and instruction at Texas A & M University in College Station, Texas. He is married to Candra Huckaby, a special education teacher at Nacogdoches Independent School District. He and Candra live in Nacogdoches.

The style manual used in this thesis is A Manual For Authors of Mathematical Papers published by the American Mathematical Society.

This thesis was prepared by Chad Alan Huckaby using L<sup>A</sup>T<sub>E</sub>X.