### Stephen F. Austin State University

# SFA ScholarWorks

**Faculty Publications** 

**Biology** 

2010

# Batrachochytrium dendrobatidis Detected in Amphibians from National Forests in Eastern Texas, USA

Daniel Saenz

Southern Research Station, U.S.D.A. Forest Service, Nacogdoches, TX 75965

Brendan T. Kavanagh

Southern Research Station, U.S.D.A. Forest Service, Nacogdoches, TX 75965

Matthew Kwiatkowski

Stephen F Austin State University, kwiatkowm@sfasu.edu

Follow this and additional works at: https://scholarworks.sfasu.edu/biology



Part of the Animal Sciences Commons, and the Biology Commons

Tell us how this article helped you.

### **Repository Citation**

Saenz, Daniel; Kavanagh, Brendan T.; and Kwiatkowski, Matthew, "Batrachochytrium dendrobatidis Detected in Amphibians from National Forests in Eastern Texas, USA" (2010). Faculty Publications. 144. https://scholarworks.sfasu.edu/biology/144

This Article is brought to you for free and open access by the Biology at SFA ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SFA ScholarWorks. For more information, please contact cdsscholarworks@sfasu.edu.

- Daszak, P., L. Berger, A. A. Cunningham, A. D. Hyatt, D. E. Green, and R. Speare. 1999. Emerging infectious diseases and amphibian population declines. Emerg. Infect. Dis. 5:735–748.
- DAVIDSON, E. W., M. PARRIS, J. P. COLLINS, J. E. LONGCORE, A. P. PESSIER, and J. L. BRUNNER. 2003. Pathengenicity and transmission of chytridiomycosis in tiger salamanders (*Ambystoma tigrinum*). Copeia 2003:601–607.
- GREEN, D. E., K. A. CONVERSE, and A. K. SCHRADER. 2002. Epizootiology of sixty-four amphibian mortality events in the USA, 1996-2001. Ann. NY Acad. Sci. 969:323–339.
- HANSELMANN, R., A. RODRÍGUEZ, M. LAMPO, L. FAJARDO-RAMOS, A. ALONSO AGUIRRE, A. MARM KILPATRICK, J. PAUL RODRÍGUEZ, and P. DASZAK. 2004. Presence of an emerging pathogen of amphibians in introduced bullfrogs *Rana catesbeiana* in Venezuela. Biol. Conserv. 120:115–119.
- Longcore, J. E., A. P. Pessier, and D. K. Nichols. 1999. *Batrachochytrium dendrobatidis*, gen. et. sp, nov., a chytrid pathogenic to amphibians. Mycologia. 91:219–227.
- NICHOLS, D. K., E. W. LAMIRANDE, A. P. PESSIER, and J. E. LONGCORE. 2001. Experimental transmission of cutaneous chytridiomycosis in dendrobatid frogs. J. Wildl. Dis. 37:1–11.
- Pallister, J., A. Gould, D. Harrison, A. D. Hyatt, J. Jancovich, and H. Heine. 2007. Development of real-time PCR assays for the detection and differentiation of Australian and European ranaviruses. J. Fish Dis. 30:427–438.
- Piotrowski, J. S., S. L. Annis, and J. E. Longcore. 2004. Physiology of *Batrachochytrium dendrobatidis*, a chytrid pathogen of amphibians. Mycologia. 96:9–15.
- RAFFEL, T. R., J. R. DILLARD, and P. J. HUDSON. 2006. Evidence for leech-bourne transmission of amphibian *Ichthyophonus* sp. J. Parasitol. 92:1256–1264.
- ROTHERMEL, B. B., S. C. WALLS, J. C. MITCHELL, C. K. DODD JR, L. K. IRWIN, D. E. GREEN, V. M. VAZQUEZ, J. W. PETRANKA, and D. J. STEVENSON. 2008. Widespread occurrence of the amphibian chytrid fungus *Batrachochytrium dendrobatidis* in the southeastern USA. Dis. Aquat. Org. 82:3–18.
- TIMPE, E. K., S. P. GRAHAM, R. W. GAGLIARDO, R. L. HILL, and M. G. Levy. 2008. Occurrence of the fungal pathogen *Batrachochytrium dendrobatidis* in Georgia's amphibian populations. Herpetol. Rev. 39:447–449.
- Vasquez, V., B. Rothermel, and A. P. Pessier. 2009. Experimental infections of North American plethodontid salamanders with the fungus *Batrachochytrium dendrobatidis*. Dis. Aquat. Org. 84:1–7.
- VENESKY, M. D., and F. M. Brem. 2008. Occurrence of *Batrachochytrium dendrobatidis* in southwestern Tennessee, USA. Herpetol. Rev. 39:319–320.
- WHITE, M. L. 1981. The Birmingham District: An Industrial History and Guide. Birmingham Historical Society, Birmingham, Alabama.

 ${\it Herpetological\ Review}, 2010, 41(1), 47–49.$  © 2010 by Society for the Study of Amphibians and Reptiles

## Batrachochytrium dendrobatidis Detected in Amphibians from National Forests in Eastern Texas, USA

### DANIEL SAENZ BRENDAN T. KAVANAGH

Southern Research Station, Forest Service, U.S. Department of Agriculture 506 Hayter Street, Nacogdoches, Texas 75965-3556, USA

#### and

### MATTHEW A. KWIATKOWSKI

Stephen F. Austin State University, Department of Biology 1936 North Street, Nacogdoches, Texas 75962-3050, USA

\*Corresponding author; e-mail: dsaenz@fs.fed.us

The amphibian disease chytridiomycosis, caused by the pathogenic fungus Batrachochytrium dendrobatidis (Bd, Longcore et al. 1999), is well known as a major threat to amphibians resulting in mass die-offs and population declines throughout the world (Berger et al. 1998; Blaustein and Keisecker 2002; Daszak et al. 2003; McCallum 2005; Rachowicz et al. 2006). Batrachochytrium dendrobatidis has been detected on amphibians from sites across North America (Ouellet et al. 2005; Woodhams et al. 2008) and appears to be most prevalent in the western and the northeastern United States (Longcore et al. 2007; Schlaepfer et al. 2007). Whereas infected anurans also have been found throughout the southeastern US (Green and Dodd 2007), there have been no reports of Bd from amphibians in eastern Texas, a broad area encompassing 10,000,000 ha. We sampled amphibians for the presence of Bd in four National Forests in eastern Texas (approximately 31°N latitude).

Amphibians were sampled for *Bd* from 9 January to 27 May 2009 in the Angelina, Davy Crockett, and Sabine National Forests, and the Stephen F. Austin Experimental Forest (Fig. 1). The Stephen




Fig. 1. Locations of the Angelina, Davy Crockett, Sabine National Forests, and the Stephen F. Austin Experimental Forest where six of 18 amphibian species tested positive for the amphibian chytrid fungus, *Batrachochytrium dendrobatidis*.

Table 1. Amphibian species tested for the presence of *Batrachochytrium dendrobatidis* (*Bd*) within Angelina, Davy Crockett, and Sabine National Forests, and the Stephen F. Austin Experimental Forest, Texas. Bold font indicates that *Bd* was detected in the species.

		No. animals infected/examined	No. animals infected/examined	No. animals infected/examined	No. animals infected/examined
Family	Species	Angelina National Forest	Davy Crockett National Forest	Sabine National Forest	Stephen F. Austin Experimental Forest
Ambystomatidae	Ambystoma maculatum	0/5	-	-	-
	Ambystoma opacum	0/10	0/2	0/5	0/10
	Ambystoma talpoideum	0/2	-	-	-
Plethodontidae	Eurycea quadradigitata	0/8	-	0/8	3/10
Salamandridae	Notophthalmus viridescens	0/1	-	-	0/2
Bufonidae	Anaxyrus woodhousii	1/7	0/10	-	0/1
	Incilius nebulifer	0/3	0/8	0/1	-
Hylidae	Acris crepitans	1/10	1/10	-	-
	Hyla cinerea	0/10	0/6	-	0/10
	Hyla versicolor	0/8	0/10	-	0/8
	Pseudacris crucifer	0/10	1/10	-	0/10
	Pseudacris fouquettei	3/5	<u>-</u>	-	2/10
Microhylidae	Gastrophryne carolinensis	0/5	-	-	0/2
Ranidae	Lithobates catesbeianus	0/1	0/6	-	1/1
	Lithobates clamitans	0/2	0/4	-	0/2
	Lithobates palustris	-	-	0/1	0/4
	Lithobates sphenocephalus	0/8	0/7	-	0/10
Scaphiopodidae	Scaphiopus hurteri	-	0/1	-	-
	TOTAL	5/95 (5.3%)	2/76 (2.6%)	0/15 (0.0%)	6/80 (7.5%)

F. Austin Experimental Forest is a disjunct unit of the Angelina National Forest, administered by the Southern Research Station, US Forest Service. The dominant habitats of these areas include secondary growth Loblolly (*Pinus taeda*), Longleaf (*P. palustris*), and Shortleaf (*P. echinata*) upland pine forests and mixed deciduous bottomland forests. East Texas experiences occasional freezing temperatures, warm winter days, and extremely hot summers (Chang et al. 1996). From 1901 to 1993, the overall mean air temperatures for January and August are 8.4°C and 27.8°C, respectively (Chang et al. 1996). In winter, cold air masses often meet warm moist air pushed up from the Gulf of Mexico resulting in frequent rain events, placing the study sites in one of the wettest regions of Texas (Bomar 1995).

We searched for amphibians near ponds, streams, moist lowland areas, and upland pine forest habitat. We captured specimens by hand. Each individual was handled with a new pair of sterile nitrile gloves. We sampled for *Bd* by rubbing a sterile cotton swab on the dorsum, ventral surfaces, and feet of each frog for approximately 30 seconds, after which the animal was released at its place of capture. The swab was then immediately placed in a sterile microcentrifuge tube containing 1 ml of 70% ethanol and later sent to Pisces Molecular Laboratory (Boulder, Colorado, USA) for PCR analyses. Global positioning system (GPS) coordinates were taken at each capture site using a Garmin® GPS unit.

Overall, we sampled a total of 266 adult amphibians of 18 different species, from 8 different families (Table 1). Of these 18 species,

six had at least one individual that tested positive for Bd. Thirteen of the 266 individuals tested positive for an overall detection rate of 4.8%. The Stephen F. Austin Experimental Forest had the highest detection rate among the four areas sampled, with six of 80 (7.5%) individuals testing positive for Bd (Table 1). During sampling, no sick or dead frogs were observed at any sites.

We found similar *Bd* detection rates for anurans and caudates (4.93% and 4.76%, respectively), although the only salamanders to test positive for *Bd* were three individuals of *Eurycea qudradigitata* from the Stephen F. Austin Experimental Forest. Despite 34 Ambystomatids being sampled, none tested positive for *Bd*.

Five of the 15 (33.33%) Pseudacris fouquettei samples tested positive for Bd, which was the highest detection rate among anuran species, including species which were breeding at the same time and place where the Bd-positive Pseudacris fouquettei were found. Individuals of both P. crucifer and Lithobates sphenocephalus were collected at locations where Bd-positive P. fouquettei were found, yet no L. sphenocephalus and only one of 30 (3.33%) P. crucifer samples tested positive for Bd. However, small sample sizes make it difficult to assess whether there is actually a higher detection rates among P. fouquettei than other species.

Batrachochytrium dendrobatidis has been present in North American amphibian populations since at least the early 1960s (Ouellet et al. 2005). Yet, within the United States, *Bd* has been associated with amphibian die-offs predominantly in western states (e.g., Bradley et al. 2002; Briggs et al. 2005; Muths et al. 2003).

Although the fungus is present in the southeastern United States, to our knowledge no amphibian declines have been attributed to chytridiomycosis in this region. Several studies have detected higher rates of *Bd* infections during the fall or winter months (McDonald et al. 2005; Ouellet et al. 2005; Retallick et al. 2004), a phenomenon possibly explained by the fact that amphibian immune systems function less effectively at cooler temperatures (Carey 2000; Cooper et al. 1992; Maniero and Carey 1997). It is possible that the immune systems of amphibians inhabiting eastern Texas are more capable of resisting *Bd* infections because of the relatively warm winters and hot summers.

Also, it is possible that the fungus may never reach an epizootic state in eastern Texas because the warm air temperatures are less than optimal for *Bd* growth and infection of amphibians (Kriger and Hero 2006; Longcore et al. 1999; Retallick et al. 2004). Evidence for an enzootic state includes the relative low incidence of detection in PCR samples and the fact that no sick or dead frogs were encountered. Amphibians appeared to be abundant at all four of our study sites. Follow-up population and *Bd* sampling is needed to confirm what impacts *Bd* may be having on the National Forests in Texas.

Acknowledgments.—We thank T. Hall, T. Cotten, S. Wahlberg, and R. Adams for assisting with field sampling. Also, we thank C. Adams, D. Laurencio, K. Floyd, and an anonymous reviewer for constructive comments on an earlier draft of this manuscript.

#### LITERATURE CITED

- Berger, L., R. Speare, P. Daszak, D. E. Green, A. A. Cunningham, C. L. Goggin, R. Slocombe, M. A. Ragan, A. D. Hyatt, K. R. McDonald, H. B. Hines, K. R. Lips, G. Martantelli, and H. Parkes. 1998. Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. Proc. Nat. Acad. Sci. 95:9031–9036.
- BLAUSTEIN, A. R. AND J. M. KEISECKER. 2002. Complexity in conservation: lessons from the global decline of amphibian populations. Ecol. Lett. 5:597–608.
- Bomar, G. 1995. The climate of the Texas hill country and East Texas. *In* J. Norwine, J. R. Giardino, G. R. North, and J. B. Valdes (eds.), The Changing Climate of Texas, Predictability and Implications for Future, pp. 76–91. Geobooks, Texas A&M University, College Station, Texas.
- Bradley, G. A., P. C. Rosen, M. J. Sredl, T. R. Jones, and J. E. Long-core. 2002. Chytridiomycosis in native Arizona frogs. J. Wildl. Dis. 38:206–212.
- BRIGGS, C. J., V. T. VREDENBURG, R. A. KNAPP, AND L. J. RACHOWICZ. 2005. Investigating the population-level effects of chytridiomycosis: an emerging infectious disease of amphibians. Ecology 86:3149–3159.
- Carey, C. 2000. Infectious disease and worldwide declines of amphibian populations, with comments on emerging diseases in coral reef organisms and in humans. Enviro. Health Perspect. 108(Suppl. 1):143–150.
- CHANG, M., L. D. CLENDENEN, AND H. C. REEVES. 1996. Characteristics of a humid climate: Nacogdoches, Texas. Center for Applied Studies in Forestry, College of Forestry, Nacogdoches, Texas.
- COOPER, E. L., R. K. WRIGHT, A. E. KLEMPAU, AND C. T. SMITH. 1992. Hibernation alters the frog's immune system. Cryobiology 29:616–631.
- DASZAK, P., A. A. CUNNINGHAM, AND A. D. HYATT. 2003. Infectious disease and amphibian declines. Diversity and Distributions 9:141–150.
- Green, D. E., AND C. K. Dodd, Jr. 2007. Presence of amphibian chytrid fungus *Batrachochytrium dendrobatidis* and other amphibian pathogens

- at warm-water fish hatcheries in southeastern North America. Herpetol. Cons. Biol. 2:43–47.
- KRIGER, K. M., AND J. M. HERO. 2006. Large-scale seasonal variation in the prevalence and severity of chytridiomycosis. J. Zool. (Lond.) 271:352–359.
- LONGCORE, J. E., A. P. PESSIER, AND D. K. NICHOLS. 1999. Batrachochytrium dendrobatidis gen. et sp. nov., a chytrid pathogenic to amphibians. Mycologia 91:219–227.
- Longcore, J. R., J. E. Longcore, A. P. Pessier, and W. A. Halteman. 2007. Chytridiomycosis widespread in anurans of northeastern United States. J. Wildl. Manage. 71:435-444.
- Maniero, G. D., and C. Carey. 1997. Changes in selected aspects of immune function in the leopard frog, *Rana pipiens*, associated with exposure to cold. J. Comp. Phys. B 167:256–263.
- McCallum, H. 2005. Inclusion of chytridiomycosis as the agent in widespread frog declines. Conserv. Biol. 19:1421–1430.
- McDonald, K. R., D. Méndez, R. Müller, A. B. Freeman, and R. Speare. 2005. Decline in the prevalence of chytridiomycosis in upland frog populations in North Queensland, Australia. Pacific Conserv. Biol. 11:114–120.
- MUTHS, E., P. S. CORN, A. P. PESSIER, AND D. E. GREEN. 2003. Evidence for disease-related decline in Colorado. Biol. Conserv. 110:357–365.
- OUELLET, M., I. MIKAELIAN, B. D. PAULI, AND J. RODRIGUE. 2005. Historical evidence of widespread chytrid infection in North American amphibian populations. Conserv. Biol. 19:1431–1440.
- RACHOWICZ, L. R., R. A. KNAPP, J. A. T. MORGAN, M. J. STICE, V. T. VRENDENBURG, J. M. PARKER, AND C. J. BRIGGS. 2006. Emerging infectious disease as a proximate cause of amphibian mass mortality. Ecology 87:1671–1683.
- RETALLICK, R. W. R., H. McCallum, and R. Speare. 2004. Endemic infection of the amphibian chytrid fungus in a frog community post-decline. PLoS 2:1965–1971.
- Schlaepfer, M. A., M. J. Sredl, P. C. Rosen, and M. J. Ryan. 2007. High prevalence of *Batrachochytrium dendrobatidis* in wild population of lowland leopard frogs *Rana yavapaiensis* in Arizona. Ecohealth 4:421-427.
- WOODHAMS, D. C., A. D. HYATT, D. G. BOYLE, AND L. A. ROLLINS-SMITH. 2008. The northern leopard frog *Rana pipiens* is a widespread reservoir species harboring *Batrachochytrium dendrobatidis* in North America. Herpetol. Rev. 39:66–68.