

# Water Soluble Cationic Porphyrin Sensor for Detection of Hg<sup>2+</sup>, Pb<sup>2+</sup>, Cd<sup>2+</sup>, and Cu<sup>2+</sup>



STEPHEN F. AUSTIN  
STATE UNIVERSITY

Department of Chemistry  
and Biochemistry



**Christopher Orr, Miranda Uherek, Matibur Zamadar\***

## ABSTRACT

Here we report the sensing properties of the aqueous solution of *meso*-tetra(N-methyl-4-pyridyl)porphine tetrachloride (**1**) for simultaneous detection of toxic metal ions by using UV-vis spectroscopy. Cationic porphyrin **1** displayed different electronic absorptions in UV-vis region upon interacting with Hg<sup>2+</sup>, Pb<sup>2+</sup>, Cd<sup>2+</sup>, and Cu<sup>2+</sup> ions in neutral water solution at room temperature. Quite interestingly, the porphyrin **1** showed that it can function as a single optical chemical sensor and/or metal ion receptor capable of detecting two or more toxic metal ions, particularly, Hg<sup>2+</sup>, Pb<sup>2+</sup>, and Cd<sup>2+</sup> ions coexisting in a water sample. Porphyrin **1** in an aqueous solution provides a unique UV-vis sensing system for the determination of Cd<sup>2+</sup> in the presence of larger metal ions such as Hg<sup>2+</sup>, or Pb<sup>2+</sup>. Finally, the examination of the sensing properties of **1** demonstrated that it can operate as a Cu<sup>2+</sup> ion selective sensor via metal displacement from the **1**-Hg<sup>2+</sup>, **1**-Pb<sup>2+</sup>, and **1**-Cd<sup>2+</sup>.






## INTRODUCTION

- Toxic metal ions are dangerous to human health and environment.
- Determination of metal ions, particularly, Hg<sup>2+</sup>, Pb<sup>2+</sup>, Cd<sup>2+</sup>, and Cu<sup>2+</sup> simultaneously by using an inexpensive, water soluble porphyrin based optical sensor has always been a great challenge.
- In addition, the preparation of porphyrin-based optical sensors often requires complex organic syntheses, which are very costly, and time consuming.
- Furthermore, the porphyrin-based sensors are water insoluble, which limits the detection of metal ions in aquatic environment.


## PURPOSE

- To find a simple, inexpensive, water soluble, and commercially available porphyrin
- a single optical chemical sensor capable of detecting any or all Hg<sup>2+</sup>, Pb<sup>2+</sup>, Cu<sup>2+</sup>, and Cd<sup>2+</sup> ions simultaneously

## RESULTS



Cd<sup>2+</sup> ions are detected in the presence of larger metal ions such as Pb<sup>2+</sup>



Cu<sup>2+</sup> ions are detected via metal displacement from the 1-Pb<sup>2+</sup>.

## FUTURE PLAN

- Preparation of solid sensor **1**
- Determination of toxic metal ions in aqueous solution using solid sensor **1**

## CONCLUSIONS

- We found water soluble, inexpensive, and commercially available cationic porphyrin **1** produced different electronic absorptions in UV-vis region upon interacting with Hg<sup>2+</sup>, Pb<sup>2+</sup>, Cd<sup>2+</sup>, and Cu<sup>2+</sup> ions.
- The porphyrin **1** showed the ability to detect multiple metal ions, particularly Hg<sup>2+</sup>, Pb<sup>2+</sup>, and Cd<sup>2+</sup> in aqueous solution.
- The porphyrin **1** displayed the ability to determine Cd<sup>2+</sup> ions more easily when other metal ions with larger ionic radii are present.
- The porphyrin **1** detected Cu<sup>2+</sup> via metal displacement from the **1**-Hg<sup>2+</sup>, **1**-Pb<sup>2+</sup>, and **1**-Cd<sup>2+</sup>.

## ACKNOWLEDGEMENTS

- Michael A. Janusa, Chair and Professor of Chemistry and Biochemistry at SFASU
- The Welch Foundation Departmental Grant (AN-0008)
- RCA grant at the Stephen F. Austin State University Research Enhancement Program